arXiv:math/0312337v2 [math.GT] 17 Dec 2003

KIRBY ELEMENTS AND QUANTUM INVARIANTS

ALEXIS VIRELIZIER

Abstract. We define the notion of a Kirby element of a ribbon category $\mathcal{C}$ (not necessarily semisimple). Kirby elements lead to 3-manifolds invariants. We characterize a set of Kirby elements (in terms of the structural morphisms of a Hopf algebra in $\mathcal{C}$) which is sufficiently large to recover the quantum invariants of 3-manifolds of Reshetikhin-Turaev, of Hennings-Kauffman-Radford, and of Lyubashenko. The cases of a semisimple ribbon category and of a category of representations are explored in details.

Contents

Introduction 1
1. Ribbon categories and coends 3
2. Kirby elements of a ribbon category 9
3. The case of semisimple ribbon categories 15
4. The case of categories of representations 25
5. A non-unimodular example 41
Appendix A. Traces on ribbon Hopf algebras 43
References 44

Introduction

During the last decade, deep connections between low-dimensional topology and the purely algebraic theory of quantum groups (or, more generally, of braided categories) were highlighted. In particular, this led to a new class of 3-manifolds invariants, called *quantum invariants*, defined in several ways.

The aim of the present paper is to give a as general as possible method of constructing quantum invariants of 3-manifolds starting from a ribbon category or a ribbon Hopf algebra. With this formalism, we recover the 3-manifolds invariants of Reshetikhin-Turaev [RT91, Tur94], of Hennings-Kauffman-Radford [KR95, Hen96], and of Lyubashenko [Lyu95a] when these are well-defined.

Let $\Bbbk$ be a field and $\mathcal{C}$ be a $\Bbbk$-linear ribbon category (not necessarily semisimple). Under some technical assumption, namely the existence of a coend $A \in \mathrm{Ob}(\mathcal{C})$ of the functor $(X, Y) \in \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \mapsto X^* \otimes Y \in \mathcal{C}$, a scalar $\tau_{\mathcal{C}}(L; \alpha)$ can be associated to any framed link L in S^3 and any morphism $\alpha \in \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, A)$, see [Lyu95a]. Recall, see [Lyu95b], that the object A of $\mathcal{C}$ is then a Hopf algebra in the category $\mathcal{C}$.

By a *Kirby element* of $\mathcal{C}$, we shall mean a morphism $\alpha \in \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, A)$ such that $\tau_{\mathcal{C}}(L; \alpha)$ is invariant under isotopies of L and under 2-handle slides. By using

---

Date: February 1, 2008.

2000 *Mathematics Subject Classification.* 57M27,18D10,81R50.

the Kirby theorem [Kir78], we have that if α is a Kirby element of $\mathcal{C}$ such that $\tau_\mathcal{C}(\bigcirc^{\pm 1};\alpha) \neq 0$, then $\tau_\mathcal{C}(L;\alpha)$ can be normalized to an invariant $\tau_\mathcal{C}(M_L;\alpha)$ of 3-manifolds, where $\bigcirc^{\pm 1}$ is the unknot with framing ±1 and M_L denotes the 3-manifold obtained from S^3 by surgery along L.

In general, determining all the Kirby elements of $\mathcal{C}$ is a quite difficult problem. In this paper we characterize, in terms of the structure maps of the categorical Hopf algebra A, a set $\mathcal{I}(\mathcal{C})$ made of Kirby elements of $\mathcal{C}$ which is sufficiently large to contain the Kirby elements corresponding to the known quantum invariants.

If the categorical Hopf algebra A admits a two-sided integral $\lambda : \mathbb{1} \to A$, then λ belongs to $\mathcal{I}(\mathcal{C})$ and the corresponding invariant $\tau_\mathcal{C}(M;\lambda)$ is the Lyubashenko invariant [Lyu95a].

When $\mathcal{C}$ is semisimple, we give necessary conditions for being in $\mathcal{I}(\mathcal{C})$ and we show that there exist (even in the non-modular case) elements of $\mathcal{I}(\mathcal{C})$ corresponding to the Reshetikhin-Turaev invariants [RT91, Tur94] computed from finitely semisimple full ribbon subcategories of $\mathcal{C}$. Note that these elements are not in general two-sided integrals.

More generally, we show that $\mathcal{I}(\mathcal{C})$ contains the Kirby elements leading to the invariants obtained from $\mathcal{I}(\mathcal{B})$, where $\mathcal{B}$ is a finitely semisimple full ribbon subcategory of the semisimple quotient of $\mathcal{C}$. We verify that, in general, there exist Kirby elements in $\mathcal{I}(\mathcal{C})$ which are not of this last form. This means that the semisimplification process "lacks" some invariants.

Let H be a finite-dimensional ribbon Hopf algebra. Suppose that $\mathcal{C}$ is the category rep_H of finite-dimensional left H-modules. We describe $\mathcal{I}(H) = \mathcal{I}(\mathrm{rep}_H)$ in purely algebraic terms. One of the interest of such a description is to avoid the representation theory of H (which may be of wild type, see [Ben98]).

If H is unimodular, then $1 \in \mathcal{I}(H)$ and the corresponding invariant $\tau_{(H,1)}$ is the Hennings-Kauffman-Radford invariant [KR95, Hen96]. More generally, and even if H is not unimodular, we show that the invariant $\tau_{(H,z)}$ of 3-manifolds corresponding to $z \in \mathcal{I}(H)$ can be computed by using the Kauffman-Radford algorithm.

If $\mathcal{V}$ is a set of simple left H-modules which makes $(H,\mathcal{V})$ a premodular Hopf algebra, then there exists $z_\mathcal{V} \in \mathcal{I}(H)$ such that $\tau_{(H,z_\mathcal{V})}$ is the Reshetikhin-Turaev invariant computed from $(H,\mathcal{V})$, which can then be computed by using the Kauffman-Radford algorithm.

When H is semisimple and $\Bbbk$ is of characteristic 0, we show that the Hennings-Kauffman-Radford invariant (computed from H) and the Reshetikhin-Turaev invariant (computed from rep_H) are simultaneously well-defined and coincide (even in the non-modular case). In the modular case, this was first shown in [Ker97].

We explicitly determine $\mathcal{I}(H)$ for a family of non-unimodular ribbon Hopf algebras which contains Sweedler's Hopf algebra.

As an algebraic application, the operators involved in the description of $\mathcal{I}(H) = \mathcal{I}(\mathrm{rep}_H)$ in algebraic terms allows us to parameterize all the traces on a finite-dimensional ribbon Hopf algebra H. When H is unimodular, we recover the parameterization given in [Rad94b, Hen96].

The paper is organized as follows. In Section 1, we review ribbon categories and coends. In Section 2, we define and study Kirby elements. We focus, in Section 3, on the case of semisimple ribbon categories and, in Section 4, on the case of categories of representations of ribbon Hopf algebras. In Section 5, we treat an example in detail. Finally, in Appendix A, we study traces on ribbon Hopf algebras.

Acknowledgements. I thank A. Bruguières for helpful discussions, in particular for questions concerning category theory.

1. Ribbon categories and coends

In this section, we review some basic definitions concerning ribbon categories and coends. Throughout this paper, we let $\Bbbk$ be a field.

1.1. **Ribbon categories.** Let $\mathcal{C}$ be a strict monoidal category with unit object $\mathbb{1}$ (note that every monoidal category is equivalent to a strict monoidal category in a canonical way, see, [Lan98]). A *left duality* in $\mathcal{C}$ associates to any object $U \in \mathcal{C}$ an object $U^* \in \mathcal{C}$ and two morphisms $\mathrm{ev}_U : U^* \otimes U \to \mathbb{1}$ and $\mathrm{coev}_U : \mathbb{1} \to U \otimes U^*$ such that

$$(\mathrm{id}_U \otimes \mathrm{ev}_U)(\mathrm{coev}_U \otimes \mathrm{id}_U) = \mathrm{id}_U \tag{1.1}$$

$$(\mathrm{ev}_U \otimes \mathrm{id}_{U^*})(\mathrm{id}_{U^*} \otimes \mathrm{coev}_U) = \mathrm{id}_{U^*}. \tag{1.2}$$

We can (and we always do) impose that $\mathbb{1}^* = \mathbb{1}$, $\mathrm{ev}_{\mathbb{1}} = \mathrm{id}_{\mathbb{1}}$ and $\mathrm{coev}_{\mathbb{1}} = \mathrm{id}_{\mathbb{1}}$.

By a *braided category* we shall mean a monoidal category $\mathcal{C}$ with left duality and endowed with a system $\{c_{U,V} : U \otimes V \to V \otimes U\}_{U,V \in \mathcal{C}}$ of invertible morphisms (the *braiding*) satisfying the following three conditions:

$$c_{U',V'}(f \otimes g) = (g \otimes f)\, c_{U,V}, \tag{1.3}$$

$$c_{U\otimes V,W} = (c_{U,W} \otimes \mathrm{id}_V)(\mathrm{id}_U \otimes c_{V,W}), \tag{1.4}$$

$$c_{U,V\otimes W} = (\mathrm{id}_V \otimes c_{U,W})(c_{U,V} \otimes \mathrm{id}_W), \tag{1.5}$$

for all objects $U, V, W \in \mathcal{C}$ and all morphisms $f : U \to U'$, $g : V \to V'$ in $\mathcal{C}$. Note that, by applying (1.4) to $U = V = \mathbb{1}$ and (1.5) to $V = W = \mathbb{1}$ and using the invertibility of $c_{U,\mathbb{1}}$ and $c_{\mathbb{1},U}$, we obtain that $c_{U,\mathbb{1}} = c_{\mathbb{1},U} = \mathrm{id}_U$ for any object $U \in \mathcal{C}$.

A *ribbon category* is a braided category $\mathcal{C}$ endowed with a family of invertible morphisms $\{\theta_U : U \to U\}_{U\in\mathcal{C}}$ (the *twist*) satisfying the following conditions:

$$\theta_V f = f\, \theta_U; \tag{1.6}$$

$$(\theta_U \otimes \mathrm{id}_{U^*})\mathrm{coev}_U = (\mathrm{id}_U \otimes \theta_{U^*})\mathrm{coev}_U; \tag{1.7}$$

$$\theta_{U\otimes V} = c_{V,U}\, c_{U,V}\, (\theta_U \otimes \theta_V); \tag{1.8}$$

for all objects $U, V \in \mathcal{C}$ and all morphism $f : U \to V$ in $\mathcal{C}$. It follows from (1.8) that $\theta_{\mathbb{1}} = \mathrm{id}_{\mathbb{1}}$.

A ribbon category $\mathcal{C}$ canonically has a right duality by associating to any object $U \in \mathcal{C}$ its left dual $U^* \in \mathcal{C}$ and two morphisms $\widetilde{\mathrm{ev}}_U : U \otimes U^* \to \mathbb{1}$ and $\widetilde{\mathrm{coev}}_U : \mathbb{1} \to U^* \otimes U$ defined by

$$\widetilde{\mathrm{ev}}_U = \mathrm{ev}_U c_{U,U^*}(\theta_U \otimes \mathrm{id}_{U^*}); \tag{1.9}$$

$$\widetilde{\mathrm{coev}}_U = (\mathrm{id}_{U^*} \otimes \theta_U^{-1})(c_{U^*,U})^{-1}\mathrm{coev}_U. \tag{1.10}$$

Note that we have $\widetilde{\mathrm{ev}}_{\mathbb{1}} = \mathrm{id}_{\mathbb{1}}$ and $\widetilde{\mathrm{coev}}_{\mathbb{1}} = \mathrm{id}_{\mathbb{1}}$.

The *dual* morphism $f^*: V^* \to U^*$ of a morphism $f: U \to V$ in a ribbon category $\mathcal{C}$ is defined by

$$\begin{aligned} f^* &= (\mathrm{ev}_V \otimes \mathrm{id}_{U^*})(\mathrm{id}_{V^*} \otimes f \otimes \mathrm{id}_{U^*})(\mathrm{id}_{V^*} \otimes \mathrm{coev}_U) \\ &= (\mathrm{id}_{U^*} \otimes \widetilde{\mathrm{ev}}_V)(\mathrm{id}_{U^*} \otimes f \otimes \mathrm{id}_{V^*})(\widetilde{\mathrm{coev}}_U \otimes \mathrm{id}_{V^*}). \end{aligned} \tag{1.11}$$

It is well-known that $(\mathrm{id}_U)^* = \mathrm{id}_{U^*}$ and $(fg)^* = g^* f^*$ for composable morphisms f, g. Axiom (1.7) can be shown to be equivalent to $\theta_U^* = \theta_{U^*}$.

Let $\mathcal{C}$ be a ribbon category. Note that $\mathrm{End}_{\mathcal{C}}(\mathbb{1})$ is a semigroup, with composition as multiplication, which is commutative (since $\mathcal{C}$ is monoidal). The *quantum trace* of an endomorphism $f: U \to U$ of an object $U \in \mathcal{C}$ is defined by

$$\mathrm{tr}_q(f) = \widetilde{\mathrm{ev}}_U(f \otimes \mathrm{id}_{U^*})\,\mathrm{coev}_U = \mathrm{ev}_U(\mathrm{id}_{U^*} \otimes f)\,\widetilde{\mathrm{coev}}_U \in \mathrm{End}_{\mathcal{C}}(\mathbb{1}). \tag{1.12}$$

For any morphisms $u: U \to V$, $v: V \to U$ and for any endomorphisms f, g in $\mathcal{C}$, we have

$$\mathrm{tr}_q(uv) = \mathrm{tr}_q(vu), \quad \mathrm{tr}_q(f^*) = \mathrm{tr}_q(f), \quad \text{and} \quad \mathrm{tr}_q(f \otimes g) = \mathrm{tr}_q(f)\,\mathrm{tr}_q(g). \tag{1.13}$$

The *quantum dimension* of an object $U \in \mathcal{C}$ is defined by

$$\dim_q(U) = \mathrm{tr}_q(\mathrm{id}_U) = \widetilde{\mathrm{ev}}_U\,\mathrm{coev}_U = \mathrm{ev}_U\,\widetilde{\mathrm{coev}}_U \in \mathrm{End}_{\mathcal{C}}(\mathbb{1}). \tag{1.14}$$

Isomorphic objects have equal dimensions and $\dim_q(U \otimes V) = \dim_q(U)\dim_q(V)$ for any objects $U, V \in \mathcal{C}$. Note that $\dim_q(\mathbb{1}) = \mathrm{id}_{\mathbb{1}}$.

1.2. **$\Bbbk$-categories.** Let $\Bbbk$ be a field. By a *$\Bbbk$-category*, we shall mean a category for which the sets of morphisms are $\Bbbk$-spaces and the composition is $\Bbbk$-bilinear. By a *monoidal $\Bbbk$-category*, we shall mean $\Bbbk$-category endowed with a monoidal structure whose tensor product is $\Bbbk$-bilinear. Note that if $\mathcal{C}$ is a monoidal $\Bbbk$-category, then $\mathrm{End}_{\mathcal{C}}(\mathbb{1})$ is a commutative $\Bbbk$-algebra (with composition as multiplication). A monoidal $\Bbbk$-category is said to be *pure* if $\mathrm{End}_{\mathcal{C}}(\mathbb{1}) = \Bbbk$.

By a *ribbon $\Bbbk$-category*, we shall mean a pure monoidal $\Bbbk$-category endowed with a ribbon structure.

1.3. **Graphical calculus.** Let $\mathcal{C}$ be ribbon category. Any morphism in $\mathcal{C}$ can be graphically represented by a plane diagram (we use the conventions of [Tur94]). This pictorial calculus will allow us to replace algebraic arguments involving commutative diagrams by simple geometric reasoning. This is justified, e.g., in [Tur94].

A morphism $f: V \to W$ in $\mathcal{C}$ is represented by a box with two vertical arrows oriented downwards, as in Figure 1(a). Here V, W should be regarded as "colors" of the arrows and f should be regarded as a "color" of the box. More generally, a morphism $f: V_1 \otimes \cdots \otimes V_m \to W_1 \otimes \cdots \otimes W_n$ may be represented as in Figure 1(b).

We also use vertical arrows oriented upwards under the convention that the morphism sitting in a box attached to such an arrow involves not the color of the arrow but rather the dual object.

The identity endomorphism of an object $V \in \mathcal{C}$ or of its dual V^* will be represented by a vertical arrow as depicted in Figure 1(c). Note that a vertical arrow colored with $\mathbb{1}$ may be deleted from any picture without changing the morphism represented by this picture. The symbol "=" displayed in the figures denotes equality of the corresponding morphisms in $\mathcal{C}$.

The tensor product $f \otimes g$ of two morphisms f and g in $\mathcal{C}$ is represented by placing a picture of f to the left of a picture of g. A picture for the composition $g \circ f$ of



(a) $f : V \to W$ (b) Tensor product (c) The identity

Figure 1. Plane diagrams of morphisms

two (composable) morphisms g and f is obtained by putting a picture of g on the top of a picture of f and by gluing the corresponding free ends of arrows.

The braiding $c_{V,W} : V \otimes W \to W \otimes V$ and its inverse $c^{-1}_{V,W} : W \otimes V \to V \otimes W$, the twist $\theta_V : V \to V$ and its inverse $\theta_V^{-1} : V \to V$, and the duality morphisms $\mathrm{ev}_V : V^* \otimes V \to \mathbb{1}$, $\mathrm{coev}_V : \mathbb{1} \to V \otimes V^*$, $\widetilde{\mathrm{ev}}_V : V \otimes V^* \to \mathbb{1}$, and $\widetilde{\mathrm{coev}}_V : \mathbb{1} \to V^* \otimes V$ are represented as in Figures 2(a), 2(b), and 2(c) respectively. The quantum trace of an endomorphism $f : V \to V$ in $\mathcal{C}$ and the quantum dimension of an object $V \in \mathcal{C}$ may be depicted as in Figure 2(d).

1.4. **Negligible morphisms.** Let $\mathcal{C}$ be a ribbon $\mathbb{k}$-category. A morphism $f \in \mathrm{Hom}_{\mathcal{C}}(X, Y)$ is said to be *negligible* if $\mathrm{tr}_q(gf) = 0$ for all $g \in \mathrm{Hom}_{\mathcal{C}}(Y, X)$. Denote by $\mathrm{Negl}_{\mathcal{C}}(X, Y)$ the $\mathbb{k}$-subspace of $\mathrm{Hom}_{\mathcal{C}}(X, Y)$ formed by the negligible morphisms.

It is important to note that $\mathrm{Negl}_{\mathcal{C}}$ is a two-sided $\otimes$-ideal of $\mathcal{C}$. This means that the composition (resp. the tensor product) of a negligible morphism with any other morphism is negligible.

Note that since $\mathrm{End}(\mathbb{1}) = \mathbb{k}$, a morphism $f : \mathbb{1} \to X$ is negligible if and only if $gf = 0$ for all morphism $g : X \to \mathbb{1}$.

1.5. **Dinatural transformations and coends.** Recall that to each category $\mathcal{C}$ we associate the *opposite category* $\mathcal{C}^{\mathrm{op}}$ in the following way: the objects of $\mathcal{C}^{\mathrm{op}}$ are the objects of $\mathcal{C}$ and the morphisms of $\mathcal{C}^{\mathrm{op}}$ are morphisms f^{op}, in one-one correspondence $f \mapsto f^{\mathrm{op}}$ with the morphisms in $\mathcal{C}$. For each morphism $f : U \to V$ of $\mathcal{C}$, the domain and codomain of the corresponding f^{op} are as in $f^{\mathrm{op}} : V \to U$ (the direction is reversed). The composite $f^{\mathrm{op}} g^{\mathrm{op}} = (gf)^{\mathrm{op}}$ is defined in $\mathcal{C}^{\mathrm{op}}$ exactly when the composite gf is defined in $\mathcal{C}$. This makes $\mathcal{C}^{\mathrm{op}}$ a category.

Let $\mathcal{C}$ and $\mathcal{B}$ be two categories. A *dinatural transformation* between a functor $F : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \to \mathcal{B}$ and an object $B \in \mathcal{B}$ is a function d which assigns to each object $X \in \mathcal{C}$ a morphism $d_X : F(X, X) \to B$ of $\mathcal{B}$ in such a way that the diagram

$$\begin{tikzcd}
F(Y, X) \arrow[r, "F(\mathrm{id}_Y, f)"] \arrow[d, "F(f, \mathrm{id}_X)"'] & F(Y, Y) \arrow[d, "d_Y"] \\
F(X, X) \arrow[r, "d_X"] & B
\end{tikzcd}$$

commutes for every morphism $f : X \to Y$ in $\mathcal{C}$.

A *coend* of the functor F is a pair (A, i) consisting of an object A of $\mathcal{B}$ and a dinatural transformation i from F to A which is *universal* among the dinatural transformation from F to a constant, that is, with the property that, to every



(a) Braiding



(b) Twist


ev_V =
coev_V =
$\widetilde{\mathrm{ev}}_V$ =
$\widetilde{\mathrm{coev}}_V$ =
V

(c) Duality morphisms



(d) Trace and dimension

FIGURE 2.

dinatural transformation d from F to B, there exists a unique morphism $r : A \to B$ such that, for all object $X \in \mathcal{C}$,

$$d_X = r \circ i_X. \tag{1.15}$$

By using the factorization property (1.15), it is easy to verify that if (A, i) and (A', i') are two coends of F, then they are isomorphic in the sense that there exists an isomorphism $I : A \to A'$ in $\mathcal{B}$ such that $i'_X = I \circ i_X$ for all object $X \in \mathcal{C}$.

1.6. **Categorical Hopf algebras from coends.** Let $\mathcal{C}$ be a ribbon category. Consider the functor $F : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \to \mathcal{C}$ defined by

$$F(X, Y) = X^* \otimes Y \quad \text{and} \quad F(f^{\mathrm{op}}, g) = f^* \otimes g \tag{1.16}$$

for all objects $X \in \mathcal{C}^{\mathrm{op}}$, $Y \in \mathcal{C}$ and all morphisms f, g in $\mathcal{C}$.

Suppose that the functor F admits a coend (A, i). Then the object A has a structure of a Hopf algebra in the category $\mathcal{C}$ (see [Lyu95b]). This means that there exist morphisms $m_A : A \otimes A \to \mathbb{1}$, $\eta_A : \mathbb{1} \to A$, $\Delta_A : A \to A \otimes A$, $\varepsilon_A : A \to \mathbb{1}$, and $S_A : A \to A$, which verify the same axioms as those of a Hopf algebra except the usual flip is replaced by the braiding $c_{A,A} : A \otimes A \to A \otimes A$. By using the factorization property (1.15) and the naturality of the braiding and twist of $\mathcal{C}$, these structural morphisms are defined as follows:

$$\Delta_A \circ i_X = \Big(X^* \otimes X \xrightarrow{\mathrm{id}_{X^*} \otimes \mathrm{coev}_X \otimes \mathrm{id}_X} X^* \otimes X \otimes X^* \otimes X \xrightarrow{i_X \otimes i_X} A \otimes A\Big), \tag{1.17}$$

$$\varepsilon_A \circ i_X = \mathrm{ev}_X, \tag{1.18}$$

$$\eta_A : \mathbb{1} = \mathbb{1}^* \otimes \mathbb{1} \xrightarrow{i_{\mathbb{1}}} A, \tag{1.19}$$

$$\begin{aligned} m_A \circ (i_X \otimes i_Y) = \Big(X^* \otimes X \otimes Y^* \otimes Y &\xrightarrow{c_{X,Y^* \otimes Y}} X^* \otimes Y^* \otimes Y \otimes X \\ &\simeq (Y \otimes X)^* \otimes (Y \otimes X) \xrightarrow{i_{Y \otimes X}} A\Big), \end{aligned} \tag{1.20}$$

$$\begin{aligned} S_A \circ i_X = \Big(X^* \otimes X \xrightarrow{c_{X^*,X}} X \otimes X^* &\xrightarrow{\mathrm{id}_X \otimes \theta_{X^*}} X \otimes X^* \\ &\simeq X^{**} \otimes X^* \xrightarrow{i_{X^*}} A\Big). \end{aligned} \tag{1.21}$$

Here X and Y are any objects of $\mathcal{C}$. The defining relations (1.17)-(1.21) are graphically represented in Figure 3.



(a) $\Delta_A : A \to A \otimes A$ (b) $\varepsilon_A : A \to 1$ (c) $\eta_A : \mathbb{1} \to A$



(d) $m_A : A \otimes A \to A$ (e) $S_A : A \to A$

FIGURE 3. Structural morphisms of A

It can be shown that the antipode $S_A : A \to A$ is an isomorphism and that $S_A^2 = \theta_A$ (see [Lyu95b]). Moreover, as for Hopf algebras, the antipode is anti-(co)multiplicative, that is,

$$S_A m_A = m_A(S_A \otimes S_A)c_{A,A}, \qquad S_A\eta_A = \eta_A, \tag{1.22}$$

$$\Delta_A S_A = c_{A,A}(S_A \otimes S_A)\Delta_A, \qquad \varepsilon_A S_A = \varepsilon_A. \tag{1.23}$$

The Hopf algebra A posses a Hopf pairing $\omega_A : A \otimes A \to \mathbb{1}$ (see [Lyu95a]) defined by

$$\omega_A \circ (i_X \otimes i_Y) = (\mathrm{ev}_X \otimes \mathrm{ev}_Y)(\mathrm{id}_{X^*} \otimes c_{Y^*,X} c_{X,Y^*} \otimes \mathrm{id}_Y) \tag{1.24}$$

for any objects $X, Y \in \mathcal{C}$. This pairing is said to be *non-degenerate* if $\omega_A(\mathrm{id}_A \otimes \mathrm{coev}_A) : A \to A^*$ and $\omega_A(\widetilde{\mathrm{coev}}_A \otimes \mathrm{id}_A) : A \to A^*$ are isomorphisms.

Set:

$$\Gamma_l = (\mathrm{id}_A \otimes m_A)(\Delta_A \otimes \mathrm{id}_A) : A \otimes A \to A \otimes A, \tag{1.25}$$

$$\Gamma_r = (m_A \otimes \mathrm{id}_A)(\mathrm{id}_A \otimes \Delta_A) : A \otimes A \to A \otimes A. \tag{1.26}$$

Lemma 1.1. $\Gamma_l(S_A \otimes S_A)c_{A,A} = c_{A,A}(S_A \otimes S_A)\Gamma_r$.

Proof. By using (1.3), (1.22) and (1.23), we have

$$\begin{aligned}
\Gamma_l(S_A \otimes S_A)c_{A,A} &= (\mathrm{id}_A \otimes m_A)(\Delta_A S_A \otimes S_A)c_{A,A}\\
&= (\mathrm{id}_A \otimes m_A)(c_{A,A}(S_A \otimes S_A)\Delta_A \otimes S_A)c_{A,A}\\
&= (S_A \otimes m_A(S_A \otimes S_A))(c_{A,A}\Delta_A \otimes \mathrm{id}_A)c_{A,A}\\
&= (S_A \otimes S_A m_A)(\mathrm{id}_A \otimes c_{A,A}^{-1})(c_{A,A} \otimes \mathrm{id}_A)(\Delta_A \otimes \mathrm{id}_A)c_{A,A}.
\end{aligned}$$

Then, using (1.3) and (1.4)-(1.5), we get that

$$\begin{aligned}
&\Gamma_l(S_A \otimes S_A)c_{A,A}\\
&\quad= (S_A \otimes S_A m_A)(\mathrm{id}_A \otimes c_{A,A}^{-1})(c_{A,A} \otimes \mathrm{id}_A)c_{A,A\otimes A}(\mathrm{id}_A \otimes \Delta_A)\\
&\quad= (S_A \otimes S_A m_A)(\mathrm{id}_A \otimes c_{A,A}^{-1})c_{A,A\otimes A}(\mathrm{id}_A \otimes c_{A,A})(\mathrm{id}_A \otimes \Delta_A)\\
&\quad= (S_A \otimes S_A m_A)(\mathrm{id}_A \otimes c_{A,A}^{-1})(\mathrm{id}_A \otimes c_{A,A})(c_{A,A} \otimes \mathrm{id}_A)(\mathrm{id}_A \otimes c_{A,A})(\mathrm{id}_A \otimes \Delta_A)\\
&\quad= (S_A \otimes S_A m_A)c_{A\otimes A,A}(\mathrm{id}_A \otimes \Delta_A)\\
&\quad= c_{A,A}(S_A \otimes S_A)(m_A \otimes \mathrm{id}_A)(\mathrm{id}_A \otimes \Delta_A)\\
&\quad= c_{A,A}(S_A \otimes S_A)\Gamma_r.
\end{aligned}$$

□

Corrolary 1.2. *Suppose that $\mathcal{C}$ is moreover a (monoidal) $\Bbbk$-category. Let $\alpha \in \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, A)$. If $S_A\alpha - \alpha \in \mathrm{Negl}_{\mathcal{C}}(\mathbb{1}, A)$, then the following assertions are equivalent:*

(a) *$\Gamma_l(\alpha \otimes \alpha) - \alpha \otimes \alpha : \mathbb{1} \to A \otimes A$ is negligible;*
(b) *$\Gamma_r(\alpha \otimes \alpha) - \alpha \otimes \alpha : \mathbb{1} \to A \otimes A$ is negligible.*

Moreover, if $S_A \circ \alpha = \alpha$, then $\Gamma_l(\alpha \otimes \alpha) = \alpha \otimes \alpha$ if and only if $\Gamma_r(\alpha \otimes \alpha) = \alpha \otimes \alpha$.

Proof. This is an immediate consequence of Lemma 1.1 since $\mathrm{Negl}_{\mathcal{C}}$ is a two-sided $\otimes$-ideal of $\mathcal{C}$. □

2. Kirby elements of a ribbon category

In this section, we generalize the Lyubashenko's method [Lyu95a] of constructing 3-manifolds invariants from ribbon categories.

2.1. **Special tangles.** Let n be a positive integer. By a *n-special tangle* we shall mean an oriented framed tangle $T \subset \mathbb{R}^2 \times [0,1]$ with $2n$ bottom endpoints and no top endpoints, consisting of n arc components such that the kth arc $(1 \leq k \leq n)$ joins the $(2k-1)$th and $(2k)$th bottom endpoints and is oriented out of $\mathbb{R}^2 \times [0,1]$ near the $(2k)$th bottom endpoint (and so inside $\mathbb{R}^2 \times [0,1]$ near the $(2k-1)$th bottom endpoint).

Diagrams of special tangles are drawn with blackboard framing. An example of a 3-special tangle is depicted in Figure 4(a).

Let $\mathcal{C}$ be a ribbon category. Suppose that the functor (1.16) admits a coend (A, i). Let T be a n-special tangle. For objets $X_1, \dots, X_n \in \mathcal{C}$, let $T_{(X_1,\cdots,X_n)}$ be the morphism $X_1^* \otimes X_1 \otimes \cdots \otimes X_n^* \otimes X_n \to \mathbb{1}$ in $\mathcal{C}$ graphically represented by a diagram of T where the kth component of T has been colored with the object X_k. Note that $T_{(X_1,\cdots,X_n)}$ does not depend on the choice of the diagram of T, that is, only depends on the isotopy class of T (see [Tur94]). Since the braiding and twist of $\mathcal{C}$ are natural and by using the Fubini theorem for coends (see [Lan98]), there exists a (unique) morphism $\phi_T : A^{\otimes n} \to \mathbb{1}$ such that

$$T_{(X_1,\dots,X_n)} = \phi_T \circ (i_{X_1} \otimes \cdots \otimes i_{X_n}) \tag{2.1}$$

for all objects $X_1, \dots, X_n \in \mathcal{C}$ (see Figure 4(b) for $n = 3$).


=
ϕ_T
A
A
A
i_{X_1}
i_{X_2}
i_{X_3}
X_1 X_1 X_2 X_2 X_3 X_3
X_1 X_1 X_2 X_2 X_3 X_3

(a) A 3-special tangle T

(b) $T_{(X_1,X_2,X_3)} = \phi_T \circ (i_{X_1} \otimes i_{X_2} \otimes i_{X_2})$

Figure 4.

2.2. **Kirby elements.** Let $\mathcal{C}$ be a ribbon category such that the functor (1.16) admits a coend (A, i).

Let L be a framed link in S^3 with n components. Fix an orientation for L. There always exists a (non-unique) n-special tangle T_L such that L is isotopic $T_L \circ (\cup_- \otimes \cdots \otimes \cup_-)$, where $\cup_-$ denote the cup with clockwise orientation, see Figure 6(a). For $\alpha \in \operatorname{Hom}_{\mathcal{C}}(\mathbb{1}, A)$, set

$$\tau_{\mathcal{C}}(L; \alpha) = \phi_{T_L} \circ \alpha^{\otimes n} \in \operatorname{End}_{\mathcal{C}}(\mathbb{1}),$$

where $\phi_{T_L} : A^{\otimes n} \to \mathbb{1}$ is defined as in (2.1).

Definition 2.1. By a *Kirby element* of $\mathcal{C}$, we shall mean a morphism $\alpha \in \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, A)$ such that, for any framed link L, $\tau_{\mathcal{C}}(L;\alpha)$ is well-defined and invariant under isotopies and 2-handle slides of L. A Kirby element α of $\mathcal{C}$ is said to be *normalized* if $\tau_{\mathcal{C}}(\bigcirc^{\pm 1};\alpha)$ is invertible in $\mathrm{End}_{\mathcal{C}}(\mathbb{1})$, where $\bigcirc^{\pm 1}$ denotes the unknot with framing ± 1.

Note that the unit $\eta_A : \mathbb{1} \to A$ of the categorical Hopf algebra A is a normalized Kirby element. The invariant of framed links associated η_A is the trivial one, that is, $\tau_{\mathcal{C}}(L;\eta_A) = 1$ for any framed link L.

Lemma 2.2. *Let α be a Kirby element of $\mathcal{C}$. Then $\tau_{\mathcal{C}}(L \sqcup L';\alpha) = \tau_{\mathcal{C}}(L;\alpha)\,\tau_{\mathcal{C}}(L';\alpha)$ for any framed link L and L', where $L \sqcup L'$ denotes the disjoint union of L and L'.*

Proof. Let T_L (resp. $T_{L'}$) be a special tangle such that L (resp. L') is isotopic $T_L \circ (\cup_- \otimes \cdots \otimes \cup_-)$ (resp. $T_{L'} \circ (\cup_- \otimes \cdots \otimes \cup_-)$). The tangle $T = T_L \otimes T_{L'}$ is then special and such that the disjoint union $L \sqcup L'$ is isotopic $T \circ (\cup_- \otimes \cdots \otimes \cup_-)$. Therefore $\phi_T = \phi_{T_L} \otimes \phi_{T_{L'}}$ and so $\tau_{\mathcal{C}}(L \sqcup L';\alpha) = \tau_{\mathcal{C}}(L;\alpha)\,\tau_{\mathcal{C}}(L';\alpha)$. □

Let $\Theta_\pm : A \to \mathbb{1}$ be the morphism defined by

$$\Theta_\pm \circ i_X = \mathrm{ev}_X(\mathrm{id}_{X^*} \otimes \theta_X^{\pm 1}), \tag{2.2}$$

where X is any object of $\mathcal{C}$ (see Figure 5). Note that if α be a Kirby element of $\mathcal{C}$, then $\tau_{\mathcal{C}}(\bigcirc^{\pm 1};\alpha) = \Theta_\pm \alpha$.



Figure 5.

Recall (see [Lic97]) that every closed, connected, and oriented 3-manifold can be obtained from S^3 by surgery along a framed link $L \subset S^3$. In this paper, all considered 3-manifolds are supposed to be closed, connected, and oriented. For any framed link L in S^3, we will denote by M_L the 3-manifold obtained from S^3 by surgery along L, by n_L the number of components of L, and by $b_-(L)$ the number of negative eigenvalues of the linking matrix of L.

Normalized Kirby elements are of special interest due to the following proposition.

Proposition 2.3. *Let α be a normalized Kirby element of $\mathcal{C}$. Then*

$$\tau_{\mathcal{C}}(M_L;\alpha) = (\Theta_+\alpha)^{b_-(L)-n_L}\,(\Theta_-\alpha)^{-b_-(L)}\,\tau_{\mathcal{C}}(L;\alpha)$$

is an invariant of 3-manifolds. Moreover $\tau_{\mathcal{C}}(M \# M';\alpha) = \tau_{\mathcal{C}}(M;\alpha)\,\tau_{\mathcal{C}}(M';\alpha)$ for any 3-manifold M and M'.

Proof. The fact that $\tau_{\mathcal{C}}(M_L;\alpha)$ is an invariant of 3-manifolds follows from Kirby theorem [Kir78]. Indeed $\tau_{\mathcal{C}}(L;\alpha)$, $b_-(L)$ and n_L are invariant under 2-handle slides and $\tau_{\mathcal{C}}(\bigcirc^{\pm 1} \sqcup L;\alpha) = (\Theta_\pm\alpha)\,\tau_{\mathcal{C}}(L;\alpha)$ by Lemma 2.2, $b_-(\bigcirc^{1} \sqcup L) = b_-(L)$, $b_-(\bigcirc^{-1} \sqcup L) = b_-(L) + 1$, and $n_{\bigcirc^{\pm 1} \sqcup L} = n_L + 1$.

Let L and L' be framed links in S^3. The disjoint union $L \sqcup L'$ is then a framed link in S^3 such that $M_{L\sqcup L'} \simeq M_L \# M_{L'}$. Then the multiplicativity of $\tau_{\mathcal{C}}(M;\alpha)$ with respect to the connected sum of 3-manifolds follows from Lemma 2.2 and the equalities $b_-(L \sqcup L') = b_-(L) + b_-(L')$ and $n_{L\sqcup L'} = n_L + n_{L'}$. □

Remarks 2.4. **[a].** For any normalized Kirby element α of $\mathcal{C}$, we have that

$$\tau_{\mathcal{C}}(S^3;\alpha) = 1 \quad \text{and} \quad \tau_{\mathcal{C}}(S^1\times S^2;\alpha) = (\Theta_+\alpha)^{-1}\,\varepsilon_A\alpha.$$

[b]. The invariant of 3-manifolds associated to the unit $\eta_A : \mathbb{1} \to A$ of the categorical Hopf algebra A (which is a normalized Kirby element) is the trivial one, that is, $\tau_{\mathcal{C}}(M;\eta_A) = 1$ for any 3-manifold M.

In general, determining when a morphism $A^{\otimes n} \to \mathbb{1}$ is of the form ϕ_T for some special n-tangle T is a quite difficult problem. Hence does so the problem of determining all the (normalized) Kirby elements of $\mathcal{C}$. In the next section, we characterize a class of (normalized) Kirby elements of $\mathcal{C}$ by means of the structural morphisms of the categorical Hopf algebra A. This class will be shown to be sufficiently large to contain the Lyubashenko invariant (which is a categorical version of the Hennings-Kauffman-Radford invariant) and the Reshetikhin-Turaev invariant (computed from a semisimple quotient of $\mathcal{C}$) when these are well-defined.

2.3. A class of Kirby elements.

Let $\mathcal{C}$ be a ribbon $\Bbbk$-category such that the functor (1.16) admits a coend (A, i).

Recall the notion of negligible morphisms (see Section 1.4). Set:

$$\begin{aligned}\mathcal{I}(\mathcal{C}) = \{\alpha \in \mathrm{Hom}_{\mathcal{C}}(\mathbb{1},A) \mid S_A\alpha - \alpha \in \mathrm{Negl}_{\mathcal{C}}(\mathbb{1},A) \quad &\text{and} \\ \Gamma_l(\alpha\otimes\alpha) - \alpha\otimes\alpha \in \mathrm{Negl}_{\mathcal{C}}(\mathbb{1}, A\otimes A)\},& \\ \mathcal{I}(\mathcal{C})^{\mathrm{norm}} = \{\alpha \in \mathcal{I}(\mathcal{C}) \mid \Theta_+\alpha \neq 0 \quad \text{and} \quad \Theta_-\alpha \neq 0\},&\end{aligned}$$

where $\Gamma_l : A\otimes A \to A\otimes A$ and $\Theta_\pm : A \to \mathbb{1}$ are defined in (1.25) and (2.2).

Note that, in the above definitions, the morphism Γ_l can be replaced by the morphism Γ_r which is defined in (1.26) (see Corollary 1.2).

Remark that the sets $\mathcal{I}(\mathcal{C})$ and $\mathcal{I}(\mathcal{C})^{\mathrm{norm}}$ always contain a non-zero element, namely the unit $\eta_A : \mathbb{1} \to A$. Moreover,

$$\Bbbk\mathcal{I}(\mathcal{C}) + \mathrm{Negl}_{\mathcal{C}}(\mathbb{1},A) \subset \mathcal{I}(\mathcal{C}) \text{ and } \Bbbk^*\mathcal{I}(\mathcal{C})^{\mathrm{norm}} + \mathrm{Negl}_{\mathcal{C}}(\mathbb{1},A) \subset \mathcal{I}(\mathcal{C})^{\mathrm{norm}}, \tag{2.3}$$

since $\mathrm{Negl}_{\mathcal{C}}$ is a two-sided $\otimes$-ideal of $\mathcal{C}$ and $\mathrm{Negl}_{\mathcal{C}}(\mathbb{1},\mathbb{1}) = 0$ (because $\mathrm{End}_{\mathcal{C}}(\mathbb{1}) = \Bbbk$ is a field).

Theorem 2.5. (a) *The elements of $\mathcal{I}(\mathcal{C})$ are Kirby elements of $\mathcal{C}$.*
(b) *The elements of $\mathcal{I}(\mathcal{C})^{\mathrm{norm}}$ are normalized Kirby elements of $\mathcal{C}$.*

Remarks 2.6. **[a].** It follows from Proposition 2.3 that any $\alpha \in \mathcal{I}(\mathcal{C})^{\mathrm{norm}}$ leads to a 3-manifolds invariant $\tau_{\mathcal{C}}(M;\alpha)$ with values in $\mathrm{End}_{\mathcal{C}}(\mathbb{1}) = \Bbbk$. Recall that this invariant is multiplicative with respect to the connected sum of 3-manifolds and verifies

$$\tau_{\mathcal{C}}(S^3;\alpha) = 1, \quad \tau_{\mathcal{C}}(S^1\times S^2;\alpha) = (\Theta_+\alpha)^{-1}\,\varepsilon_A\alpha, \quad \tau_{\mathcal{C}}(M;\eta_A) = 1,$$

for any 3-manifold M.

[b]. Let $\alpha \in \mathcal{I}(\mathcal{C})^{\mathrm{norm}}$, $n \in \mathrm{Negl}_{\mathcal{C}}(\mathbb{1},A)$, and $k \in \Bbbk^*$. Then

$$k\alpha + n \in \mathcal{I}(\mathcal{C})^{\mathrm{norm}} \quad \text{and} \quad \tau_{\mathcal{C}}(M; k\alpha+n) = \tau_{\mathcal{C}}(M;\alpha)$$

for any 3-manifold M. This follows from (2.3) and the choice of the normalization in the definition of $\tau_{\mathcal{C}}(M;\alpha)$ (see Proposition 2.3).

[c]. Suppose that $\mathcal{C}$ is 3-modular (see [Lyu95a]). Then A admits a (non-zero) two-sided integral $\lambda \in \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, A)$, that is,

$$m_A(\lambda \otimes \mathrm{id}_A) = \eta_A \varepsilon_A = m_A(\mathrm{id}_A \otimes \lambda).$$

Moreover $\lambda \in \mathcal{I}(\mathcal{C})^{\mathrm{norm}}$ and $\tau_{\mathcal{C}}(M;\lambda)$ is the Lyubashenko invariant of 3-manifolds. Note that when $\mathcal{C}$ admits split idempotents, then λ is unique (up to scalar multiple), see [BKLT00].

The condition that A posses a (non-zero) two-sided integral is quite limitative (for example, when $\mathcal{C}$ is the category rep_H of representations of a finite-dimensional Hopf algebra H, this implies that H must be unimodular). In Section 5, we give an example of non-unimodular ribbon Hopf algebra H and of an element $\alpha \in \mathcal{I}(\mathrm{rep}_H)$ which is not a two-sided integral and leads to a non-trivial invariant.

[d]. When $\mathcal{C}$ is finitely semisimple, we show in Section 3.3 (see Corollary 3.7) that there exists $\alpha_{\mathcal{C}} \in \mathcal{I}(\mathcal{C})$ corresponding to the Reshetikhin-Turaev invariant defined with $\mathcal{C}$. Note that $\alpha_{\mathcal{C}} \in \mathcal{I}(\mathcal{C})$ is not in general a two-sided integral.

More generally, we show in Section 3.5 (see Corollary 3.9) that $\mathcal{I}(\mathcal{C})$ contains elements corresponding to the Reshetikhin-Turaev invariants defined with finitely semisimple full ribbon subcategories of the ribbon semisimple quotient of $\mathcal{C}$.

Proof of Theorem 2.5. Let us prove Part (a). Fix $\alpha \in \mathcal{I}(\mathcal{C})$. Let $L = L_1 \cup \cdots \cup L_n$ be a framed link. Firstly, since $S_A\alpha - \alpha \in \mathrm{Negl}_{\mathcal{C}}(\mathbb{1}, A)$, then $\tau_{\mathcal{C}}(L;\alpha)$ does not depend on choice of T_L neither the orientation of L and is an isotopic invariant of the framed link L. Indeed, this is proved in the case $S_A\alpha = \alpha$ in [Lyu95a, Proposition 5.2.1]. The same arguments work when $S_A\alpha - \alpha \in \mathrm{Negl}_{\mathcal{C}}(\mathbb{1}, A)$ since $\mathrm{Negl}_{\mathcal{C}}$ is a two-sided $\otimes$-ideal of $\mathcal{C}$.

Let us show that $\tau_{\mathcal{C}}(L;\alpha)$ is invariant under 2-handle slides. Choose an orientation for L. Without loss of generality, we can suppose that the component L_1 slides over L_2. Let L'_2 be a copy of L_2 (following the framing) and set $L' = (L_1 \# L'_2) \cup L_2 \cup \cdots \cup L_n$. We have to show that $\tau_{\mathcal{C}}(L';\alpha) = \tau_{\mathcal{C}}(L;\alpha)$. Let T_L be a n-special tangle such that L is isotopic $T_L \circ (\cup_- \otimes \cdots \otimes \cup_-)$, where the ith cup (with clockwise orientation) corresponds to the component L_i, see Figure 6(a). Let $\Delta_2(T_L)$ be the $(2n+2, 0)$-tangle obtained by copying the 2nd component of T_L (following the framing) in such a way that the endpoints of the new component are between the 2nd and 3th bottom endpoints and between the 4th and 5th bottom endpoints of T_L. A n-special tangle T' such that L' is isotopic $T' \circ (\cup_- \otimes \cdots \otimes \cup_-)$ (where the ith cup corresponds to the ith component of L') can be constructed from $\Delta_2(T_L)$ as in Figure 6(b). For example, if T_L is the 3-special tangle depicted in Figure 6(c), then let T' be the 3-special tangle of Figure 6(d).
By the equalities of Figure 2.3 where $X_1, \ldots, X_n$ are any objects of $\mathcal{C}$, and by the uniqueness of the factorisation through a coend, we get that

$$\phi_{T'} = \phi_{T_L} \circ (\Gamma_l \otimes \mathrm{id}_{A^{\otimes(n-2)}}).$$

Therefore, since $\Gamma_l(\alpha \otimes \alpha) - \alpha \otimes \alpha \in \mathrm{Negl}_{\mathcal{C}}(\mathbb{1}, A \otimes A)$, we get that

$$\tau_{\mathcal{C}}(L';\alpha) = \phi_{T'}\alpha^{\otimes n} = \phi_{T_L}(\mathrm{id}_{A^{\otimes 2}} \otimes \alpha^{\otimes(n-2)})\Gamma_l(\alpha \otimes \alpha) = \phi_{T_L}\alpha^{\otimes n} = \tau_{\mathcal{C}}(L;\alpha).$$

To show Part (b), it suffices to remark that $\tau_{\mathcal{C}}(\bigcirc^{\pm 1};\alpha) \in \mathrm{End}_{\mathcal{C}}(\mathbb{1})$ is invertible since $\tau_{\mathcal{C}}(\bigcirc^{\pm 1};\alpha) = \Theta_{\pm}\alpha \neq 0$ and $\mathrm{End}_{\mathcal{C}}(\mathbb{1}) = \mathbb{k}$ is a field. $\square$



(a) $L \sim T_L \circ (\cup_- \otimes \cdots \otimes \cup_-)$ (b) T'



(c) T_L (d) T'

Figure 6.

2.4. **Kirby elements via functors.** Let us see that Kirby elements can be "pulled back" via ribbon functors.

Let $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$ be ribbon $\Bbbk$-categories. Suppose that the functor (1.16) of $\mathcal{A}$ admits a coend (A, i) and that the functor (1.16) of $\mathcal{B}$ admits a coend (B, j). Let $\pi : \mathcal{A} \to \mathcal{C}$ and $\iota : \mathcal{B} \to \mathcal{C}$ be ribbon functors. Since A and B are categorical Hopf algebras and π and ι are ribbon functors, the objects $\pi(A)$ and $\iota(B)$ are Hopf algebras in $\mathcal{C}$ (with structure maps induced by π and ι respectively).

Proposition 2.7. *Suppose that π is surjective, ι is full and faithful, and that there exists a Hopf algebra morphism $\varphi : \iota(B) \to \pi(A)$ such that $\pi(i_X) = \varphi \circ \iota(j_Y)$ for all objects $X \in \mathcal{A}$ and $Y \in \mathcal{B}$ with $\pi(X) = \iota(Y)$. Let $\beta \in \mathrm{Hom}_{\mathcal{B}}(\mathbb{1}, B)$ and $\alpha \in \mathrm{Hom}_{\mathcal{A}}(\mathbb{1}, A)$ such that $\pi(\alpha) = \varphi \circ \iota(\beta)$.*

- (a) *If $\beta \in \mathcal{I}(\mathcal{B})$, then $\alpha \in \mathcal{I}(\mathcal{A})$ and $\tau_{\mathcal{A}}(L; \alpha) = \tau_{\mathcal{B}}(L; \beta)$ for any framed link L.*
- (b) *If $\beta \in \mathcal{I}(\mathcal{B})^{\mathrm{norm}}$, then $\alpha \in \mathcal{I}(\mathcal{A})^{\mathrm{norm}}$ and $\tau_{\mathcal{A}}(M; \alpha) = \tau_{\mathcal{B}}(M; \beta)$ for any 3-manifold M.*

Proof. Let us prove Part (a). Suppose that $\beta \in \mathcal{I}(\mathcal{B})$. Since the structure maps of $\pi(A)$ and $\iota(B)$ are induced by π and ι from those of A and B respectively, and since $\varphi : \iota(B) \to \pi(A)$ is a Hopf algebra morphism, we have:

$$\begin{aligned}\pi(S_A\alpha - \alpha) &= (S_{\pi(A)} - \mathrm{id}_{\pi(A)})\pi(\alpha) = (S_{\pi(A)} - \mathrm{id}_{\pi(A)})\varphi\iota(\beta)\\ &= \varphi(S_{\iota(B)} - \mathrm{id}_{\iota(B)})\iota(\beta) = \varphi\iota(S_B\beta - \beta)\end{aligned}$$



FIGURE 7.

and

$$\begin{aligned}
\pi(\Gamma_l^A(\alpha\otimes\alpha)-\alpha\otimes\alpha) &= (\Gamma_l^{\pi(A)}-\mathrm{id}_{\pi(A)^{\otimes 2}})(\pi(\alpha)\otimes\pi(\alpha))\\
&= (\Gamma_l^{\pi(A)}-\mathrm{id}_{\pi(A)^{\otimes 2}})\varphi\iota(\beta\otimes\beta)\\
&= \varphi(\Gamma_l^{\iota(B)}-\mathrm{id}_{\iota(B)^{\otimes 2}})\iota(\beta\otimes\beta)\\
&= \varphi\iota(\Gamma_l^{B}(\beta\otimes\beta)-\beta\otimes\beta).
\end{aligned}$$

Now, since $S_B\beta-\beta$ and $\Gamma_l^B(\beta\otimes\beta)-\beta\otimes\beta$ are negligible in $\mathcal{B}$, ι is full, and $\mathrm{tr}_q^{\mathcal{B}}=\mathrm{tr}_q^{\mathcal{C}}\circ\iota$, we get that $\pi(S_A\alpha-\alpha)$ and $\pi(\Gamma_l^A(\alpha\otimes\alpha)-\alpha\otimes\alpha)$ are negligible in $\mathcal{C}$. Hence, since π is surjective and $\mathrm{tr}_q^{\mathcal{C}}\circ\pi=\mathrm{tr}_q^{\mathcal{A}}$, the morphisms $S_A\alpha-\alpha$ and $\Gamma_l^A(\alpha\otimes\alpha)-\alpha\otimes\alpha$ are negligible in $\mathcal{A}$, that is, $\alpha\in\mathcal{I}(\mathcal{A})$.

Let $L=L_1\cup\cdots\cup L_n$ be a framed link in S^3. Let T_L be a n-special tangle such that L is isotopic $T_L\circ(\cup_-\otimes\cdots\otimes\cup_-)$, where the ith cup (with clockwise orientation) corresponds to the component L_i. Let $Y_1,\dots,Y_n$ be any objects of $\mathcal{B}$. Since π is surjective, there exists objets $X_1,\dots,X_n$ of $\mathcal{A}$ such that $\pi(X_k)=\iota(Y_k)$. Recall that, by assumption, $\pi(i_{X_k})=\varphi\iota(j_{Y_k})$. Since ι is full and the domain and codomain of the morphism $\pi(\phi_{T_L}^{\mathcal{A}})\varphi^{\otimes n}$ of $\mathcal{C}$ are $\iota(B^{\otimes n})$ and $\mathbb{1}=\iota(\mathbb{1})$ respectively, there exists a morphism $\xi : B^{\otimes n} \to \mathbb{1}$ in $\mathcal{B}$ such that $\iota(\xi) = \pi(\phi^{\mathcal{A}}_{T_L})\varphi^{\otimes n}$. Then

$$\begin{aligned}\iota\big(\phi^{\mathcal{B}}_{T_L} \circ (j_{Y_1} \otimes \cdots \otimes j_{Y_n})\big) &= \iota(T^{\mathcal{B}}_{L(Y_1,\dots,X_n)})\\ &= T^{\mathcal{C}}_{L(\iota(Y_1),\dots,\iota(Y_n))}\\ &= T^{\mathcal{C}}_{L(\pi(X_1),\dots,\pi(X_n))}\\ &= \pi(T^{\mathcal{A}}_{L(X_1,\dots,X_n)})\\ &= \pi(\phi^{\mathcal{A}}_{T_L})(\pi(i_{X_1}) \otimes \cdots \otimes \pi(i_{X_n}))\\ &= \pi(\phi^{\mathcal{A}}_{T_L})(\varphi\iota(j_{Y_1}) \otimes \cdots \otimes \varphi\iota(j_{Y_n}))\\ &= \pi(\phi^{\mathcal{A}}_{T_L})\varphi^{\otimes n} \circ \iota(j_{Y_1} \otimes \cdots \otimes j_{Y_n})\\ &= \iota\big(\xi \circ (j_{Y_1} \otimes \cdots \otimes j_{Y_n})\big).\end{aligned}$$

Therefore, since ι is faithful, $\phi^{\mathcal{B}}_{T_L} \circ (j_{Y_1} \otimes \cdots \otimes j_{Y_n}) = \xi \circ (j_{Y_1} \otimes \cdots \otimes j_{Y_n})$ and so, by the uniqueness of the factorization through a coend, we get that $\phi^{\mathcal{B}}_{T_L} = \xi$, that is, $\iota(\phi^{\mathcal{B}}_{T_L}) = \pi(\phi^{\mathcal{A}}_{T_L})\varphi^{\otimes n}$. Hence, since the maps $\mathrm{End}_{\mathcal{A}}(\mathbb{1}) = \Bbbk \to \mathrm{End}_{\mathcal{C}}(\mathbb{1}) = \Bbbk$ and $\mathrm{End}_{\mathcal{B}}(\mathbb{1}) = \Bbbk \to \mathrm{End}_{\mathcal{C}}(\mathbb{1}) = \Bbbk$ induced by π and ι respectively are the identity of $\Bbbk$, we have that

$$\begin{aligned}\tau_{\mathcal{C}}(L;\alpha) = \pi(\tau_{\mathcal{C}}(L;\alpha)) &= \pi(\phi^{\mathcal{A}}_{T_L}\alpha^{\otimes n})\\ &= \pi(\phi^{\mathcal{A}}_{T_L})\varphi^{\otimes n}\iota(\beta^{\otimes n})\\ &= \iota(\phi^{\mathcal{B}}_{T_L}\beta^{\otimes n}) = \iota(\tau_{\mathcal{B}}(L;\beta)) = \tau_{\mathcal{B}}(L;\beta).\end{aligned}$$

Let us prove Part (b). Suppose that $\beta \in \mathcal{I}(\mathcal{B})^{\mathrm{norm}}$. Let Y be any objet of $\mathcal{B}$. Since π is surjective, there exists an object X of $\mathcal{A}$ such that $\pi(X) = \iota(Y)$. Recall that, by assumption, $\pi(i_X) = \varphi \circ \iota(j_Y)$. Since ι is full and the domain and codomain of the morphism $\pi(\Theta^{\mathcal{A}}_{\pm})\varphi$ of $\mathcal{C}$ are $\iota(B)$ and $\mathbb{1} = \iota(\mathbb{1})$ respectively, there exists a morphism $\varsigma_{\pm} : B \to \mathbb{1}$ in $\mathcal{B}$ such that $\iota(\varsigma_{\pm}) = \pi(\Theta^{\mathcal{A}}_{\pm})\varphi$. We have that

$$\begin{aligned}\iota(\varsigma_{\pm} \circ \iota(j_Y)) = \pi(\Theta^{\mathcal{A}}_{\pm})\varphi \circ \iota(j_Y) &= \pi(\Theta^{\mathcal{A}}_{\pm} i_X) = \pi(\mathrm{ev}^{\mathcal{A}}_X(\mathrm{id}_{X^*} \otimes \theta^{\mathcal{A}\pm 1}_X))\\ &= \mathrm{ev}^{\mathcal{C}}_{\pi(X)}(\mathrm{id}_{\pi(X)^*} \otimes \theta^{\mathcal{C}\pm 1}_{\pi(X)}) = \iota(\mathrm{ev}^{\mathcal{B}}_Y)(\iota(\mathrm{id}_{Y^*}) \otimes \iota(\theta^{\mathcal{B}}_Y)^{\pm 1})\\ &= \iota(\mathrm{ev}^{\mathcal{B}}_Y(\mathrm{id}_{Y^*} \otimes \theta^{\mathcal{B}\pm 1}_Y)) = \iota(\Theta^{\mathcal{B}}_{\pm} \circ j_Y).\end{aligned}$$

Therefore, since ι is faithful, $\Theta^{\mathcal{B}}_{\pm} \circ j_Y = \varsigma_{\pm} \circ \iota(j_Y)$ and so, by the uniqueness of the factorization through a coend, we get that $\Theta^{\mathcal{B}}_{\pm} = \varsigma_{\pm}$, that is, $\iota(\Theta^{\mathcal{B}}_{\pm}) = \pi(\Theta^{\mathcal{A}}_{\pm})\varphi$. Then

$$\Theta^{\mathcal{C}}_{\pm}\alpha = \pi(\Theta^{\mathcal{A}}_{\pm}\alpha) = \pi(\Theta^{\mathcal{A}}_{\pm})\varphi\iota(\beta) = \iota(\Theta^{\mathcal{B}}_{\pm}\beta) = \Theta^{\mathcal{B}}_{\pm}\beta.$$

Hence, since $\Theta^{\mathcal{B}}_{\pm}\beta \neq 0$ and by Part (a), we get that $\alpha \in \mathcal{I}(\mathcal{A})^{\mathrm{norm}}$ and $\tau_{\mathcal{A}}(M;\alpha) = \tau_{\mathcal{B}}(M;\beta)$ for any 3-manifold M. □

3. The case of semisimple ribbon categories

In this section, we focus on the case of semisimple categories $\mathcal{C}$. We give necessary conditions for being in $\mathcal{I}(\mathcal{C})$. In particular, we show that there exist (even in the non-modular case) elements of $\mathcal{I}(\mathcal{C})$ corresponding to Reshetikhin-Turaev invariants computed with finitely semisimple ribbon subcategories of $\mathcal{C}$. Moreover, we study Kirby elements coming from semisimplification of ribbon categories.

3.1. **Semisimple categories.** Recall that a category $\mathcal{C}$ *admits (finite) direct sums* if, for any finite set of objects $X_1, \dots, X_n$ of $\mathcal{C}$, there exists an object X and morphisms $p_i : X \to X_i$ such that, for any object Y and morphisms $f_i : Y \to X_i$, there is a unique morphism $f : Y \to X$ with $p_i \circ f = f_i$ for all i. The object X is then unique up to isomorphism. We write $X = \oplus_i X_i$ and $f = \oplus_i f_i$.

A $\Bbbk$-category is *abelian* if it admits (finite) direct sums, every morphism has a kernel and a cokernel, every monomorphism is the kernel of its cokernel, every epimorphism is the cokernel of its kernel, and every morphism is expressible as the composite of an epimorphism followed by a monomorphism. In particular, an abelian category admits a null object (which is unique up to isomorphism). Note that a morphism of an abelian $\Bbbk$-category which is both a monomorphism and an epimorphism is an isomorphism.

Let $\mathcal{C}$ be an abelian $\Bbbk$-category. A non-null object U of $\mathcal{C}$ is *simple* if every non-zero monomorphism $V \to U$ is an isomorphism, and every non-zero epimorphism $U \to V$ is an isomorphism. Any non-zero morphism between simple objects is an isomorphism. An object U of $\mathcal{C}$ is *scalar* if $\mathrm{End}_{\mathcal{C}}(V) = \Bbbk$. Note that if $\Bbbk$ is algebraically closed, then every simple object is scalar. An object of $\mathcal{C}$ is *indecomposable* if it cannot be written as a direct sum of two non-null objects. Note that every scalar or simple object is indecomposable.

By a *semisimple* $\Bbbk$*-category*, we shall mean an abelian $\Bbbk$-category for which every object is a (finite) direct sum of simple objects. By a *finitely semisimple* $\Bbbk$*-category*, we shall mean a semisimple $\Bbbk$-category which has finitely many isomorphism classes of simple objects. Note that in a semisimple $\Bbbk$-category, every scalar or indecomposable object is simple.

In a semisimple ribbon $\Bbbk$-category $\mathcal{C}$, any negligible morphism is null. Therefore, for every object X, Y of $\mathcal{C}$, the pairing

$$\mathrm{Hom}_{\mathcal{C}}(X,Y) \otimes \mathrm{Hom}_{\mathcal{C}}(Y,X) \to \Bbbk \quad f \otimes g \mapsto \mathrm{tr}_q(gf) \tag{3.1}$$

is non-degenerate. Note that this implies that the quantum dimension of a scalar object of $\mathcal{C}$ is invertible.

Lemma 3.1. *Let $\mathcal{C}$ be a finitely semisimple ribbon $\Bbbk$-category whose simple objects are scalar. Let Λ be a (finite) set of representatives of isomorphism classes of simple objects of $\mathcal{C}$. Fix an object X of $\mathcal{C}$. For any $\lambda \in \Lambda$, set*

$$n_\lambda = \dim_{\Bbbk} \mathrm{Hom}_{\mathcal{C}}(\lambda, X) = \dim_{\Bbbk} \mathrm{Hom}_{\mathcal{C}}(X, \lambda)$$

and let $\{f_i^\lambda \,|\, 1 \leq i \leq n_\lambda\}$ be a basis of $\mathrm{Hom}_{\mathcal{C}}(\lambda, X)$ and $\{g_i^\lambda \,|\, 1 \leq i \leq n_\lambda\}$ be a basis of $\mathrm{Hom}_{\mathcal{C}}(X, \lambda)$ such that $g_i^\lambda f_j^\lambda = \delta_{i,j}\, \mathrm{id}_\lambda$ for all $1 \leq i,j \leq n_\lambda$ (such basis exist since the pairing (3.1) *is non-degenerate). Then*

$$\mathrm{id}_X = \sum_{\lambda \in \Lambda} \sum_{1 \leq i \leq n_\lambda} f_i^\lambda g_i^\lambda .$$

Proof. Note that since $\mathcal{C}$ is semisimple with scalar simple objects, the $\mathrm{Hom}_{\mathcal{C}}$'s $\Bbbk$-spaces are finite-dimensional. Since the category $\mathcal{C}$ is semisimple, the composition induces a $\Bbbk$-linear isomorphism $\oplus_{\lambda \in \Lambda} \mathrm{Hom}_{\mathcal{C}}(X, \lambda) \otimes \mathrm{Hom}_{\mathcal{C}}(\lambda, X) \to \mathrm{End}_{\mathcal{C}}(X)$. Therefore, for all $\lambda \in \Lambda$ and $1 \leq i,j \leq n_\lambda$, there exist $a_{\lambda,i,j} \in \Bbbk$ such that

$\mathrm{id}_X=\sum_{\lambda\in\Lambda}\sum_{1\leq i,j\leq n_\lambda} a_{\lambda,i,j}\, f_i^\lambda g_j^\lambda$. Let $\lambda\in\Lambda$ and $1\leq i,j\leq n_\lambda$. Then

$$\begin{aligned}\delta_{i,j}\,\mathrm{id}_\lambda = g_i^\lambda f_j^\lambda = g_i^\lambda\, \mathrm{id}_X\, f_j^\lambda &= \sum_{\mu\in\Lambda}\sum_{1\leq k,l\leq n_\lambda} a_{\mu,k,l}\, g_i^\lambda f_k^\mu g_l^\mu f_j^\lambda \\ &= \sum_{\mu\in\Lambda}\sum_{1\leq k,l\leq n_\lambda} a_{\mu,k,l}\,\delta_{\lambda,\mu}\,\delta_{i,k}\,\delta_{j,l}\,\mathrm{id}_\mu = a_{\lambda,i,j}\,\mathrm{id}_\lambda,\end{aligned}$$

and so $a_{\lambda,i,j}=\delta_{i,j}$. Hence $\mathrm{id}_X=\sum_{\lambda\in\Lambda}\sum_{1\leq i\leq n_\lambda} f_i^\lambda g_i^\lambda$. □

3.2. **Kirby elements of finitely semisimple ribbon categories.**

Let $\mathcal{B}$ be a finitely semisimple ribbon $\Bbbk$-category whose simple objects are scalar.

Note that under these assumptions, the $\Bbbk$-space $\mathrm{Hom}_\mathcal{B}(X,Y)$ is finite-dimensional for any objects X,Y of $\mathcal{B}$.

Denote by Λ a (finite) set of representatives of isomorphism classes of simple objects of $\mathcal{B}$. We can suppose that $\mathbb{1}\in\Lambda$. For any $\lambda\in\Lambda$, there exists a unique $\lambda^\vee\in\Lambda$ such that $\lambda^*\simeq\lambda^\vee$. The map ${}^\vee:\Lambda\to\Lambda$ is an involution and $\mathbb{1}^\vee=\mathbb{1}$. Recall that $\dim_q(\lambda)\neq 0$ for any $\lambda\in\Lambda$.

Set

$$B=\bigoplus_{\lambda\in\Lambda}\lambda^*\otimes\lambda\in\mathcal{B}. \tag{3.2}$$

In particular, there exist morphisms $p_\lambda: B\to\lambda^*\otimes\lambda$ and $q_\lambda:\lambda^*\otimes\lambda\to B$ such that $\mathrm{id}_B=\sum_{\lambda\in\Lambda} q_\lambda p_\lambda$ and $p_\lambda q_\mu=\delta_{\lambda,\mu}\,\mathrm{id}_{\lambda^*\otimes\lambda}$. Let X be an object of $\mathcal{B}$. Since $\mathcal{B}$ is semisimple, we can write $X=\oplus_{i\in I}\lambda_i$, where I is a finite set and $\lambda_i\in\Lambda$. We set

$$j_X=\sum_{i\in I} q_{\lambda_i}\circ(q_i^*\otimes p_i): X^*\otimes X\to B, \tag{3.3}$$

where $p_i: X\to\lambda_i$ and $q_i:\lambda_i\to X$ are morphisms in $\mathcal{B}$ such that $\mathrm{id}_X=\sum_{i\in I} q_ip_i$ and $p_iq_j=\delta_{i,j}\,\mathrm{id}_{\lambda_i}$. Note that j_X does not depend on the choice of such a family of morphisms. Remark that $j_\lambda=q_\lambda$ for any $\lambda\in\Lambda$. It is not difficult to verify that (B,j) is a coend of the functor (1.16) of $\mathcal{B}$. By Section 1.6, the object B is a Hopf algebra in $\mathcal{B}$.

For any $\lambda\in\Lambda$, set

$$e_\lambda=j_\lambda\widetilde{\mathrm{coev}}_\lambda:\mathbb{1}\to B \quad\text{and}\quad f_\lambda=\mathrm{ev}_\lambda p_\lambda: B\to\mathbb{1}. \tag{3.4}$$

Note that $f_\lambda e_\mu=\delta_{\lambda,\mu}\dim_q(\lambda)$ for any $\lambda,\mu\in\Lambda$.

Lemma 3.2. (a) $(e_\lambda)_{\lambda\in\Lambda}$ *is a basis of the* $\Bbbk$*-space* $\mathrm{Hom}_\mathcal{B}(\mathbb{1},B)$.

(b) $(f_\lambda)_{\lambda\in\Lambda}$ *is a basis of the* $\Bbbk$*-space* $\mathrm{Hom}_\mathcal{B}(B,\mathbb{1})$.

(c) *For any* $\lambda,\mu\in\Lambda$,

$$\begin{aligned} &S_B(e_\lambda)=e_{\lambda^\vee}, \qquad \eta_B=e_\mathbb{1}, \qquad \varepsilon_B(e_\lambda)=\dim_q(\lambda),\\ &(\mathrm{id}_B\otimes f_\mu)\Delta_B(e_\lambda)=\delta_{\lambda,\mu}e_\lambda, \qquad m_B(e_\lambda\otimes e_\mu)=\sum_{\nu\in\Lambda} N_{\lambda,\mu}^\nu e_\nu,\end{aligned}$$

where $N_{\lambda,\mu}^\nu=\dim_\Bbbk\mathrm{Hom}_\mathcal{B}(\lambda\otimes\mu,\nu)$.

Proof. Let us prove Part (a). For any $\lambda\in\Lambda$, the $\Bbbk$-space $\mathrm{Hom}_\mathcal{B}(\mathbb{1},\lambda^*\otimes\lambda)$ is one-dimensional (since λ is scalar) with basis $\widetilde{\mathrm{coev}}_\lambda$. Therefore there exists $x_\lambda\in\Bbbk$ such that $p_\lambda f=x_\lambda\widetilde{\mathrm{coev}}_\lambda$, and so $f=\mathrm{id}_Bf=\sum_{\lambda\in\Lambda}q_\lambda p_\lambda f=\sum_{\lambda\in\Lambda}x_\lambda e_\lambda$. Hence the family $(e_\lambda)_{\lambda\in\Lambda}$ generates $\mathrm{Hom}_\mathcal{B}(\mathbb{1},B)$. To show that it is free, suppose that $\sum_{\lambda\in\Lambda}x_\lambda e_\lambda=0$. Then, for any $\mu\in\Lambda$, $0=\sum_{\lambda\in\Lambda}x_\lambda f_\mu e_\lambda=\dim_q(\mu)x_\mu$ and so $x_\mu=0$ since $\dim_q(\mu)\neq 0$.



(a) $S_B(e_\lambda) = e_{\lambda^\vee}$



(b) $(\mathrm{id}_B \otimes f_\mu)\Delta_B(e_\lambda) = \delta_{\lambda,\mu} e_\lambda$

FIGURE 8.

Part (b) can be shown similarly. Let us prove Part (c). Let $\lambda, \mu \in \Lambda$. By definition, $\eta_B = j_{\mathbb{1}} = j_{\mathbb{1}} \widetilde{\mathrm{coev}}_{\mathbb{1}} = e_{\mathbb{1}}$ (since $\widetilde{\mathrm{coev}}_{\mathbb{1}} = \mathrm{id}_{\mathbb{1}}$) and $\varepsilon_B(e_\lambda) = \varepsilon_B j_\lambda \widetilde{\mathrm{coev}}_\lambda = \mathrm{ev}_\lambda \widetilde{\mathrm{coev}}_\lambda = \dim_q(\lambda)$. The equalities $S_B(e_\lambda) = e_{\lambda^\vee}$ and $(\mathrm{id}_B \otimes f_\mu)\Delta_B(e_\lambda) = \delta_{\lambda,\mu} e_\lambda$ are shown in Figure 8(a) and 8(b) respectively. Write $\lambda \otimes \mu = \oplus_{i \in I} \lambda_i$. In particular, there exits morphisms $p_i : \lambda \otimes \mu \to \lambda_i$ and $q_i : \lambda_i \to \lambda \otimes \mu$ such that $\mathrm{id}_{\lambda \otimes \mu} = \sum_{i \in I} q_i p_i$ and $p_i q_j = \delta_{i,j} \mathrm{id}_{\lambda_i}$. Recall that $j_{\lambda \otimes \mu} = \sum_{i \in I} j_{\lambda_i}(q_i^* \otimes p_i)$. For any $\nu \in \Lambda$, since $\mathrm{Hom}_{\mathcal{B}}(\lambda \otimes \mu, \nu) \cong \oplus_{i \in I} \mathrm{Hom}_{\mathcal{B}}(\lambda_i, \nu)$, we have that

$$N_{\lambda,\mu}^{\nu} = \dim_{\mathbb{k}} \mathrm{Hom}_{\mathcal{B}}(\lambda \otimes \mu, \nu) = \sum_{i \in I} \delta_{\lambda_i, \nu}.$$

Then the equality $m_B(e_\lambda \otimes e_\mu) = \sum_{\nu \in \Lambda} N_{\lambda,\mu}^{\nu} e_\nu$ is shown in Figure 9. □

Since $\mathcal{B}$ is semisimple and so its negligible morphisms are null, we have that:

$$\mathcal{I}(\mathcal{B}) = \{\alpha \in \mathrm{Hom}_{\mathcal{B}}(\mathbb{1}, A) \mid S_A \alpha = \alpha \text{ and } \Gamma_l(\alpha \otimes \alpha) = \alpha \otimes \alpha\}.$$

Lemma 3.3. *Let* $\alpha = \sum_{\lambda \in \Lambda} \alpha_\lambda e_\lambda \in \mathrm{Hom}_{\mathcal{B}}(\mathbb{1}, B)$*, where* $\alpha_\lambda \in \mathbb{k}$*. Suppose that* $\alpha \in \mathcal{I}(\mathcal{B})$*. Set* $\Lambda_\alpha = \{\lambda \in \Lambda \,|\, \alpha_\lambda \neq 0\}$*. Then* $\alpha = \alpha_{\mathbb{1}} \sum_{\lambda \in \Lambda_\alpha} \dim_q(\lambda) e_\lambda$*. Moreover, the set* Λ_α *is such that* $\Lambda_\alpha^\vee = \Lambda_\alpha$ *and* $m_B(\alpha \otimes e_\lambda) = \dim_q(\lambda)\, \alpha = m_B(e_\lambda \otimes \alpha)$ *for all* $\lambda \in \Lambda_\alpha$*.*

Proof. By Lemma 3.2(c), since $S_B(\alpha) = \alpha$, we have that $\alpha_\lambda = \alpha_{\lambda^\vee}$ for all $\lambda \in \Lambda$ and so $\Lambda_\alpha^\vee = \Lambda_\alpha$. Let $\mu, \nu \in \Lambda$. Since $\Gamma_r(\alpha \otimes \alpha) = \alpha \otimes \alpha$ and by using Lemma 3.2(c),



FIGURE 9. $\Delta_B(e_\lambda \otimes e_\mu) = \sum_{\nu\in\Lambda} N^{\nu}_{\lambda,\mu} e_\nu$

we have that

$$\begin{aligned}
\mathrm{dim}_q(\mu)\mathrm{dim}_q(\nu)\alpha_\nu\alpha_\mu &= (f_\nu \otimes f_\lambda)(\alpha\otimes\alpha) = (f_\nu\otimes f_\mu)\Gamma_r(\alpha\otimes\alpha)\\
&= \sum_{\lambda,\omega\in\Lambda} \alpha_\lambda\alpha_\omega\, (f_\nu\otimes f_\mu)\Gamma_r(e_\lambda\otimes e_\omega)\\
&= \sum_{\lambda,\omega\in\Lambda} \alpha_\lambda\alpha_\omega\, f_\nu m_B(e_\lambda\otimes(\mathrm{id}_B\otimes f_\mu)\Delta_B(e_\omega))\\
&= \sum_{\lambda\in\Lambda} \alpha_\lambda\alpha_\mu\, f_\nu m_B(e_\lambda\otimes e_\mu)\\
&= \sum_{\lambda,\omega\in\Lambda} \alpha_\lambda\alpha_\mu N^{\omega}_{\lambda,\mu}\, f_\nu e_\omega\\
&= \sum_{\lambda\in\Lambda} \alpha_\lambda\alpha_\mu N^{\nu}_{\lambda,\mu}\mathrm{dim}_q(\nu),
\end{aligned}$$

and so

$$\mathrm{dim}_q(\mu)\alpha_\mu\alpha_\nu = \alpha_\mu \sum_{\lambda\in\Lambda} N^{\nu}_{\lambda,\mu}\alpha_\lambda. \tag{3.5}$$

Note that $N^{\mathbb{1}}_{\lambda,\mu} = \mathrm{dim}_{\Bbbk}\,\mathrm{Hom}_{\mathcal{B}}(\lambda\otimes\mu,\mathbb{1}) = \mathrm{dim}_{\Bbbk}\,\mathrm{Hom}_{\mathcal{B}}(\lambda,\mu^*) = \delta_{\lambda,\mu^\vee}$ for all $\lambda,\mu\in\Lambda$. Hence (3.5) gives that $\mathrm{dim}_q(\mu)\alpha_\mu\alpha_{\mathbb{1}} = \alpha_\mu\alpha_{\mu^\vee} = \alpha_\mu^2$ and so $\alpha_\mu = \alpha_{\mathbb{1}}\mathrm{dim}_q(\mu)$

whenever $\alpha_\mu \neq 0$, that is, $\alpha = \alpha_{\mathbb{1}} \sum_{\lambda\in\Lambda_\alpha} \dim_q(\lambda) e_\lambda$. Finally, for any $\mu \in \Lambda_\alpha$,

$$\begin{aligned} m_B(\alpha\otimes e_\mu) &= \sum_{\lambda\in\Lambda} \alpha_\lambda\, m_B(e_\lambda\otimes e_\mu) \\ &= \sum_{\lambda,\nu\in\Lambda} N^\nu_{\lambda,\mu}\alpha_\lambda\, e_\nu \\ &= \sum_{\nu\in\Lambda} \dim_q(\mu)\alpha_\nu e_\nu \quad \text{by (3.5) since } \alpha_\mu\neq 0 \\ &= \dim_q(\mu)\alpha. \end{aligned}$$

Likewise, using the fact that $\Gamma_l(\alpha\otimes\alpha) = \alpha\otimes\alpha$, on gets that $m_B(e_\mu\otimes\alpha) = \dim_q(\mu)\alpha$. □

By Lemma 3.3, determining $\mathcal{I}(\mathcal{B})$ resumes to find the subsets E of Λ for which $\sum_{\lambda\in E}\dim_q(\lambda)e_\lambda$ belongs to $\mathcal{I}(\mathcal{B})$. In the next theorem, we show that among these subsets, there are those corresponding to monoidal subcategories of $\mathcal{B}$.

Theorem 3.4. *Let $\mathcal{D}$ be a full ribbon and abelian subcategory of $\mathcal{B}$. Let $\Lambda_\mathcal{D}$ be a (finite) set of representatives of isomorphism classes of simple objects of $\mathcal{D}$. We can suppose that $\Lambda_\mathcal{D}\subset\Lambda$. Then $\sum_{\lambda\in\Lambda_\mathcal{D}}\dim_q(\lambda)e_\lambda \in \mathcal{I}(\mathcal{B})$.*

Remarks 3.5. **[a].** We do not know if every element of $\mathcal{I}(\mathcal{B})$ is of this form.

[b]. In Section 3.3, we verify that $\sum_{\lambda\in\Lambda_\mathcal{D}}\dim_q(\lambda)e_\lambda$ leads to the Reshetikhin-Turaev invariant defined with $\mathcal{D}$.

[c]. When $\mathcal{B}$ is modular in the sense that the pairing $\omega_B : B\otimes B\to\mathbb{1}$ defined in (1.24) is non-degenerate or, equivalently, that the S-matrix $[\mathrm{tr}_q(c_{\mu,\lambda}\circ c_{\lambda,\mu})]_{\lambda,\mu\in\Lambda}$ is inversible, then $\sum_{\lambda\in\Lambda}\dim_q(\lambda)e_\lambda$ is a two-sided integral of B (see [Ker97]) and so belongs to $\mathcal{I}(\mathcal{B})$. Nevertheless, $\sum_{\lambda\in\Lambda}\dim_q(\lambda)e_\lambda$ is not in general a two-sided integral of B.

Proof of Theorem 3.4. Firstly, since $\Lambda_\mathcal{D}^\vee = \Lambda_\mathcal{D}$ and by Lemma 3.2(c),

$$S_B(\alpha_\mathcal{D}) = \sum_{\lambda\in\Lambda_\mathcal{D}}\dim_q(\lambda)S_B(e_\lambda) = \sum_{\lambda\in\Lambda_\mathcal{D}}\dim_q(\lambda^\vee)e_{\lambda^\vee} = \alpha_\mathcal{D}.$$

Secondly, to show that $\Gamma_r(\alpha_\mathcal{D}\otimes\alpha_\mathcal{D}) = \alpha_\mathcal{D}\otimes\alpha_\mathcal{D}$, it suffices to show that, for any $\lambda\in\Lambda_\mathcal{D}$,

$$\Gamma_r(\alpha_\mathcal{D}\otimes e_\lambda) = \alpha_\mathcal{D}\otimes e_\lambda. \tag{3.6}$$

Fix $\lambda\in\Lambda_\mathcal{D}$. Let $\mu,\nu\in\Lambda_\mathcal{D}$ and set $n_{\mu,\nu} = \dim_\Bbbk \mathrm{Hom}_\mathcal{B}(\nu\otimes\lambda,\mu)$. Since $\mathcal{D}$ is a full subcategory of $\mathcal{B}$ and the pairing

$$g\otimes f\in\mathrm{Hom}_\mathcal{B}(\nu\otimes\lambda,\mu)\otimes\mathrm{Hom}_\mathcal{B}(\mu,\nu\otimes\lambda)\mapsto\mathrm{tr}_q(gf)\in\Bbbk$$

is non-degenerate, there exist basis $\{f_i^{\mu,\nu}\,|\,1\leq i\leq n_{\mu,\nu}\}$ of $\mathrm{Hom}_\mathcal{B}(\mu,\nu\otimes\lambda)$ and $\{g_i^{\mu,\nu}\,|\,1\leq i\leq n_{\mu,\nu}\}$ of $\mathrm{Hom}_\mathcal{B}(\nu\otimes\lambda,\mu)$ such that $g_j^{\mu,\nu}f_i^{\mu,\nu} = \delta_{i,j}\,\mathrm{id}_\mu$ for all $1\leq i,j\leq n_{\mu,\nu}$. By Lemma 3.1, we have that

$$\sum_{\mu\in\Lambda}\sum_{1\leq i\leq n_{\mu,\nu}} f_i^{\mu,\nu}g_i^{\mu,\nu} = \mathrm{id}_{\nu\otimes\lambda}. \tag{3.7}$$

Let $\mu,\nu\in\Lambda_{\mathcal{D}}$. For any $1\le i\le n_{\mu,\nu}$, set

$$\begin{aligned} F_i^{\nu,\mu} &= (g_i^{\mu,\nu}\otimes\mathrm{id}_\lambda)(\mathrm{id}_\nu\otimes\widetilde{\mathrm{coev}}_\lambda):\nu\to\mu\otimes\lambda^*,\\ G_i^{\nu,\mu} &= (\mathrm{id}_\nu\otimes\widetilde{\mathrm{ev}}_\lambda)(f_i^{\mu,\nu}\otimes\mathrm{id}_\lambda):\mu\otimes\lambda^*\to\nu. \end{aligned}$$

Since $(\lambda^*,\widetilde{\mathrm{ev}}_\lambda,\widetilde{\mathrm{coev}}_\lambda)$ is a right duality for λ, we have that $\{F_i^{\nu,\mu}\,|\,1\le i\le n_{\mu,\nu}\}$ is a basis for $\mathrm{Hom}_{\mathcal{B}}(\nu,\mu\otimes\lambda^*)$ and $\{G_i^{\nu,\mu}\,|\,1\le i\le n_{\mu,\nu}\}$ is a basis for $\mathrm{Hom}_{\mathcal{B}}(\mu\otimes\lambda^*,\nu)$. For any $1\le i,j\le n_{\mu,\nu}$, since $G_j^{\nu,\mu}F_i^{\nu,\mu}\in\mathrm{End}_{\mathcal{B}}(\nu)$ and ν is scalar, we have that

$$\begin{aligned} \dim_q(\nu)\,G_j^{\nu,\mu}F_i^{\nu,\mu} &= \mathrm{tr}_q(G_j^{\nu,\mu}F_i^{\nu,\mu})\,\mathrm{id}_\nu=\mathrm{tr}_q(F_i^{\nu,\mu}G_j^{\nu,\mu})\,\mathrm{id}_\nu\\ &= \mathrm{tr}_q((g_i^{\mu,\nu}\otimes\mathrm{id}_\lambda)(\mathrm{id}_\nu\otimes\widetilde{\mathrm{coev}}_\lambda)(\mathrm{id}_\nu\otimes\widetilde{\mathrm{ev}}_\lambda)(f_j^{\mu,\nu}\otimes\mathrm{id}_\lambda))\,\mathrm{id}_\nu\\ &= \mathrm{tr}_q(g_i^{\mu,\nu}f_j^{\mu,\nu})\,\mathrm{id}_\nu=\mathrm{tr}_q(\delta_{i,j}\,\mathrm{id}_\mu)\,\mathrm{id}_\nu\\ &= \delta_{i,j}\dim_q(\mu)\,\mathrm{id}_\nu. \end{aligned}$$

Therefore, by Lemma 3.1,

$$\sum_{\nu\in\Lambda}\sum_{1\le i\le n_{\mu,\nu}}\dim_q(\nu)\,F_i^{\nu,\mu}G_i^{\nu,\mu}=\dim_q(\mu)\,\mathrm{id}_{\mu\otimes\lambda^*}. \tag{3.8}$$

Finally one gets (3.6) by the equalities depicted in Figure 10 which follow from the dinaturality of j, the definition of m_B and Δ_B, and equalities (3.7) and (3.8). □

Recall that an object X of $\mathcal{B}$ is invertible if $X^*\otimes X$ is isomorphic to $\mathbb{1}$.

Corrolary 3.6. *Suppose that every simple object of $\mathcal{B}$ is invertible or, equivalently, that $(\Lambda,\otimes,\mathbb{1})$ is a group. Then $\mathcal{I}(\mathcal{B})$ is composed by the scalar multiples of $\alpha_G=\sum_{\lambda\in G}\dim_q(\lambda)e_\lambda$, where G is any subgroup of Λ.*

Proof. Let G be a subgroup of Λ. The full abelian subcategory $\mathcal{D}$ of $\mathcal{B}$ generated by G is ribbon. Therefore, by Theorem 3.4, the element $\alpha_G=\sum_{\lambda\in G}\dim_q(\lambda)e_\lambda$ (and so its scalar multiples) belongs to $\mathcal{I}(\mathcal{B})$.

Conversely, let $\alpha=\sum_{\lambda\in\Lambda}\alpha_\lambda e_\lambda\in\mathcal{I}(\mathcal{B})$. If $\alpha=0$, then $\alpha=0\cdot\alpha_\Lambda$. Suppose that $\alpha\neq 0$. By Lemma 3.3, $\alpha=\alpha_{\mathbb{1}}\sum_{\lambda\in\Lambda_\alpha}\dim_q(\lambda)e_\lambda$ where $\Lambda_\alpha=\{\lambda\in\Lambda\,|\,\alpha_\lambda\neq 0\}$. Then $\mathbb{1}\in\Lambda_\alpha$. Since every element of Λ is invertible, we have that $N_{\lambda,\mu}^{\nu}=\delta_{\nu,\lambda\otimes\mu}$ for any $\lambda,\mu,\nu\in\Lambda$. Let $\mu,\nu\in\Lambda_\alpha$. By using (3.5), $0\neq\dim_q(\mu)\alpha_\mu\alpha_\nu=\alpha_\mu\alpha_{\nu\otimes\mu^\vee}$ and so $\alpha_{\nu\otimes\mu^\vee}\neq 0$, that is, $\nu\otimes\mu^\vee\in\Lambda_\alpha$. Hence Λ_α is a subgroup of Λ. □

3.3. On the Reshetikhin-Turaev invariant.

Let $\mathcal{B}$ be a finitely semisimple ribbon $\Bbbk$-category whose simple objects are scalar. Let Λ be a (finite) set of representatives of isomorphism classes of simple objects containing $\mathbb{1}$.

Set $\Delta_\pm=\sum_{\lambda\in\Lambda}v_\lambda^{\pm1}\dim_q(\lambda)^2\in\Bbbk$, where $v_\lambda\in\Bbbk$ is the (invertible) scalar defined by $\theta_\lambda=v_\lambda\,\mathrm{id}_\lambda$. Recall (see [Tur94, Bru00a]) that the Reshetikhin-Turaev invariant of 3-manifolds is well-defined when $\Delta_+\neq 0\neq\Delta_-$. Moreover, if L is a framed link in S^3, it is given by:

$$\mathrm{RT}_{\mathcal{B}}(M_L)=\Delta_+^{b_-(L)-n_L}\,\Delta_-^{-b_-(L)}\sum_{c\in\mathrm{Col}(L)}\Big(\prod_{j=1}^{n}\dim_q(c(L_j))\Big)F(L,c)\in\Bbbk.$$

Here $\mathrm{Col}(L)$ is the set of functions $c:\{L_1,\dots,L_n\}\to\Lambda$, where $L_1,\dots,L_n$ are the components of L, and $F(L,c)\in\mathrm{End}_{\mathcal{B}}(\mathbb{1})=\Bbbk$ is the morphism represented by a plane diagram of L where the component L_j is colored with the object $c(L_j)$.

By Theorem 3.4, $\alpha_{\mathcal{B}}=\sum_{\lambda\in\Lambda}\dim_q(\lambda)e_\lambda\in\mathcal{I}(\mathcal{B})$, where e_λ is defined as in Section 3.2.


B
Γ_r
$\alpha_\mathcal{D}$
e_λ
$= \sum_{\nu\in\Lambda_\mathcal{D}} \dim_q(\nu)$
$j_{\lambda\otimes\nu}$
$\mathrm{id}_{\lambda\otimes\nu}$
$\lambda\otimes\nu$
j_λ
λ
ν
$= \sum_{\nu\in\Lambda_\mathcal{D}} \dim_q(\nu)$
$j_{\nu\otimes\lambda}$
$\mathrm{id}_{\nu\otimes\lambda}$
$\nu\otimes\lambda$
$= \sum_{\substack{\mu,\nu\in\Lambda_\mathcal{D}\\ 1\leq i\leq n_{\mu,\nu}}} \dim_q(\nu)$
$f_i^{\mu,\nu}$
$g_i^{\mu,\nu}$
μ
j_μ
$= \sum_{\mu\in\Lambda_\mathcal{D}} \sum_{\substack{\nu\in\Lambda_\mathcal{D}\\ 1\leq i\leq n_{\mu,\nu}}} \dim_q(\nu)$
$F_i^{\nu,\mu}$
$G_i^{\nu,\mu}$
$= \sum_{\mu\in\Lambda_\mathcal{D}} \dim_q(\mu)$
$=$

FIGURE 10. $\Gamma_r(\alpha_\mathcal{D}\otimes e_\lambda) = \alpha_\mathcal{D}\otimes e_\lambda$

Corrolary 3.7. *Suppose that* $\Delta_\pm \neq 0$. *Then* $\alpha_\mathcal{B} \in \mathcal{I}(\mathcal{B})^{\mathrm{norm}}$ *and* $\tau_\mathcal{B}(M;\alpha_\mathcal{B}) = \mathrm{RT}_\mathcal{B}(M)$ *for any 3-manifold* M.

Proof. We have that

$$\Theta_\pm \alpha_\mathcal{B} = \sum_{\lambda\in\Lambda} \dim_q(\lambda)\,\mathrm{ev}_\lambda(\mathrm{id}_{\lambda^*}\otimes\theta_\lambda^{\pm1})\widetilde{\mathrm{coev}}_\lambda = \sum_{\lambda\in\Lambda} v_\lambda^{\pm1}\dim_q(\lambda)^2 = \Delta_\pm \neq 0.$$

Therefore, since $\alpha_\mathcal{B}\in\mathcal{I}(\mathcal{B})$, one gets that $\alpha_\mathcal{B}\in\mathcal{I}(\mathcal{B})^{\mathrm{norm}}$.

Let $L = L_1\cup\cdots\cup L_n$ be a framed link in S^3. Let T_L be a n-special tangle such that L is isotopic $T_L\circ(\cup_-\otimes\cdots\otimes\cup_-)$, where the ith cup (with clockwise orientation) corresponds to the component L_i. Then, for any $c \in \mathrm{Col}(L)$,

$$F(L,c) = \phi_{T_L} \circ (j_{c(L_1)}\widetilde{\mathrm{coev}}_{c(L_1)} \otimes \cdots \otimes j_{c(L_n)}\widetilde{\mathrm{coev}}_{c(L_n)}) = \phi_{T_L} \circ (e_{c(L_1)} \otimes \cdots \otimes e_{c(L_n)}).$$

Therefore

$$\begin{aligned} &\sum_{c\in \mathrm{Col}(L)} \Big(\prod_{j=1}^{n} \dim_q(c(L_j))\Big) F(L,\lambda) \\ &\quad = \sum_{c\in \mathrm{Col}(L)} \phi_{T_L} \circ \Big(\dim_q(c(L_1))\, e_{c(L_1)} \otimes \cdots \otimes \dim_q(c(L_n)) e_{c(L_n)}\Big) \\ &\quad = \phi_{T_L} \circ \Big(\sum_{\lambda_1\in\Lambda} \dim_q(\lambda_1) e_{\lambda_1} \otimes \cdots \otimes \sum_{\lambda_n\in\Lambda} \dim_q(\lambda_n) e_{\lambda_n}\Big) \\ &\quad = \phi_{T_L} \circ (\alpha_{\mathcal{B}} \otimes \cdots \otimes \alpha_{\mathcal{B}}). \end{aligned}$$

Hence

$$\begin{aligned} \mathrm{RT}_{\mathcal{B}}(M_L) &= \Delta_+^{b_-(L)-n_L}\, \Delta_-^{-b_-(L)} \sum_{\lambda\in \mathrm{Col}(L)} \Big(\prod_{j=1}^{n} \dim_q(\lambda(L_j))\Big) F(L,\lambda), \\ &= (\Theta_+ \alpha_{\mathcal{B}})^{b_-(L)-n_L}\, (\Theta_- \alpha_{\mathcal{B}})^{-b_-(L)}\, \phi_{T_L} \circ (\alpha_{\mathcal{B}} \otimes \cdots \otimes \alpha_{\mathcal{B}}) \\ &= \tau_{\mathcal{B}}(M_L; \alpha_{\mathcal{B}}). \end{aligned}$$

□

3.4. **Semisimplification of ribbon categories.** Let $\mathcal{C}$ be a ribbon $\Bbbk$-category. For any objects $X, Y \in \mathcal{C}$, recall that $\mathrm{Negl}_{\mathcal{C}}(X,Y)$ denotes the $\Bbbk$-space of negligible morphisms of $\mathcal{C}$ from X to Y (see Section 1.4). Let $\mathcal{C}^s$ be the category whose objects are the same as in $\mathcal{C}$, and whose morphisms are

$$\mathrm{Hom}_{\mathcal{C}^s}(X,Y) = \mathrm{Hom}_{\mathcal{C}}(X,Y)/\mathrm{Negl}_{\mathcal{C}}(X,Y) \tag{3.9}$$

for any objets $X, Y \in \mathcal{C}^s$. The composition, monoidal structure, braiding, twist, and duality of $\mathcal{C}^s$ are induced by those of $\mathcal{C}$.

When $\mathcal{C}$ has finite-dimensional Hom's $\Bbbk$-spaces, the category $\mathcal{C}^s$ is a semisimple ribbon $\Bbbk$-category, called the *semisimplification* of $\mathcal{C}$, and the simple objects of $\mathcal{C}^s$ are the indecomposable objects of $\mathcal{C}$ with non-zero quantum dimension, see [Bru00b].

Let $\pi : \mathcal{C} \to \mathcal{C}^s$ be the functor defined by $\pi(X) = X$ and $\pi(f) = f + \mathrm{Negl}_{\mathcal{C}}(X,Y)$ for any object X and any morphism $f : X \to Y$ in $\mathcal{C}$. This is a surjective ribbon functor. Note that π is bijective on the objects.

3.5. **Kirby elements from semisimplification.** Let $\mathcal{C}$ be a ribbon $\Bbbk$-category which admits a coend (A, i) for the functor (1.16) and whose Hom's spaces are finite-dimensional. Denote by $\mathcal{C}^s$ the semisimplification of $\mathcal{C}$ and let $\pi : \mathcal{C} \to \mathcal{C}^s$ be its associated surjective ribbon functor (see Section 3.4).

Let $\mathcal{B}$ be a full ribbon and abelian subcategory of $\mathcal{C}^s$ which admits finitely many isomorphism classes of simple objects, and whose simple objects are scalar. Let Λ be a (finite) set of representatives of isomorphism classes of simple objects of $\mathcal{B}$ containing $\mathbb{1}$. For any object X of $\mathcal{C}^s$, we denote by $\pi^{-1}(X)$ the (unique) object of $\mathcal{C}$ such that $\pi(\pi^{-1}(X)) = X$.

Let $B = \oplus_{\lambda\in\Lambda} \lambda^* \otimes \lambda$. In particular, there exist morphisms $p_\lambda : B \to \lambda^* \otimes \lambda$ and $q_\lambda : \lambda^* \otimes \lambda \to B$ of $\mathcal{B}$ such that $\mathrm{id}_B = \sum_{\lambda\in\Lambda} q_\lambda p_\lambda$ and $p_\lambda q_\mu = \delta_{\lambda,\mu}\, \mathrm{id}_{\lambda^*\otimes\lambda}$. For any object X of $\mathcal{B}$, we let $j_X : X^* \otimes X \to B$ as in (3.3). Recall that (B, j) is a coend for the functor (1.16) of $\mathcal{B}$ and that $j_\lambda = q_\lambda$ for any $\lambda \in \Lambda$ (see Section 3.2).

Since $\mathcal{B}$ is a full ribbon subcategory of $\mathcal{C}^s$ and π is a ribbon functor, the objects B and $\pi(A)$ are Hopf algebra in $\mathcal{C}^s$. Set

$$\varphi = \sum_{\lambda\in\Lambda} \pi(i_{\pi^{-1}(\lambda)})p_\lambda \in \mathrm{Hom}_{\mathcal{C}^s}(B, \pi(A)). \tag{3.10}$$

Lemma 3.8. *$\varphi : B \to \pi(A)$ is a Hopf algebra morphism such that $\pi(i_X) = \varphi j_{\pi(X)}$ for all object X of $\mathcal{C}$ with $\pi(X) \in \mathcal{B}$.*

Proof. Let X be an object of $\mathcal{C}$ such that $\pi(X) \in \mathcal{B}$. Since $\mathcal{B}$ is semisimple, we can write $\pi(X) = \oplus_{k\in K}\lambda_k$, where K is a finite set and $\lambda_k \in \Lambda$. Recall that $j_{\pi(X)} = \sum_{k\in K} q_{\lambda_k}(q_k^* \otimes p_k)$, where $p_k : \pi(X) \to \lambda_k$ and $q_k : \lambda_k \to \pi(X)$ are morphisms in $\mathcal{B}$ such that $\mathrm{id}_{\pi(X)} = \sum_{k\in K} q_k p_k$ and $p_k q_l = \delta_{k,l}\,\mathrm{id}_{\lambda_k}$. For any $k \in K$, since π surjective, there exist morphisms $P_k : X \to \pi^{-1}(\lambda_k)$ and $Q_k : \pi^{-1}(\lambda_k) \to X$ in $\mathcal{C}$ such that $\pi(P_k) = p_k$ and $\pi(Q_k) = q_k$. Then, using the dinaturality of i and since the functor π is ribbon,

$$\begin{aligned}
\varphi j_{\pi(X)} &= \sum_{\lambda\in\Lambda}\sum_{k\in K} \pi(i_{\pi^{-1}(\lambda)})p_\lambda q_{\lambda_k}(q_k^* \otimes p_k)\\
&= \sum_{\lambda\in\Lambda}\sum_{\substack{k\in K\\ \lambda_k=\lambda}} \pi(i_{\pi^{-1}(\lambda)})(q_k^* \otimes p_k)\\
&= \sum_{\lambda\in\Lambda}\sum_{\substack{k\in K\\ \lambda_k=\lambda}} \pi\big(i_{\pi^{-1}(\lambda)}(Q_k^* \otimes P_k)\big)\\
&= \sum_{\lambda\in\Lambda}\sum_{\substack{k\in K\\ \lambda_k=\lambda}} \pi\big(i_X(\mathrm{id}_{X^*} \otimes Q_k P_k)\big)\\
&= \pi(i_X)(\mathrm{id}_{\pi(X)^*} \otimes \sum_{k\in K} q_k p_k)\\
&= \pi(i_X)(\mathrm{id}_{\pi(X)^*} \otimes \mathrm{id}_{\pi(X)}) = \pi(i_X).
\end{aligned}$$

Let us verify that φ is a Hopf algebra morphism. For any objects X, Y of a ribbon category, set

$$\beta_X = (\widetilde{\mathrm{ev}}_X \otimes \mathrm{id}_{X^{**}})(\mathrm{id}_X \otimes \mathrm{coev}_{X^*}) : X \xrightarrow{\simeq} X^{**}$$

and

$$\begin{aligned}
\gamma_{X,Y} =&(\mathrm{ev}_{X^*} \otimes \mathrm{id}_{(Y\otimes X)^*})(\mathrm{id}_{X^*} \otimes \mathrm{ev}_{Y^*} \otimes \mathrm{id}_{(Y\otimes X)^*})\\
&(\mathrm{id}_{X^*\otimes Y^*} \otimes \mathrm{coev}_{Y\otimes X}) : X^* \otimes Y^* \xrightarrow{\simeq} (Y \otimes X)^*.
\end{aligned}$$

Let $\lambda, \mu \in \Lambda$. Set $U = \pi^{-1}(\lambda)$ and $V = \pi^{-1}(\mu)$. Then

$$\varepsilon_{\pi(A)}\varphi j_\lambda = \pi(\varepsilon_A i_U) = \pi(\mathrm{ev}_U) = \mathrm{ev}_\lambda = \varepsilon_B j_\lambda,$$

$$\begin{aligned}
\Delta_{\pi(A)}\varphi j_\lambda &= \pi(\Delta_A i_U)\\
&= \pi((i_U \otimes i_U)(\mathrm{id}_{U^*} \otimes \mathrm{coev}_U \otimes \mathrm{id}_U))\\
&= (\varphi j_\lambda \otimes \varphi j_\lambda)(\mathrm{id}_{\lambda^*} \otimes \mathrm{coev}_\lambda \otimes \mathrm{id}_\lambda) = (\varphi \otimes \varphi)\Delta_B j_\lambda,
\end{aligned}$$

$$\begin{aligned} m_{\pi(A)}(\varphi j_\lambda \otimes \varphi j_\mu) &= \pi(m_A(i_U \otimes i_V)) \\ &= \pi(i_{V\otimes U}(\gamma_{U,V} \otimes \mathrm{id}_{V\otimes U})(\mathrm{id}_{U^*} \otimes c_{U,V^*\otimes V})) \\ &= \varphi j_{\mu\otimes\lambda}(\gamma_{\lambda,\mu} \otimes \mathrm{id}_{\mu\otimes\lambda})(\mathrm{id}_{\lambda^*} \otimes c_{\lambda,\mu^*\otimes\mu})) \\ &= \varphi m_B(j_\lambda \otimes j_\mu), \end{aligned}$$

and

$$\begin{aligned} S_{\pi(A)}\varphi j_\lambda = \pi(S_A i_U) &= \pi(i_{U^*}(\beta_U \otimes \theta_{U^*}^{-1})c_{U^*,U}) \\ &= \varphi j_{\lambda^*}(\beta_\lambda \otimes \theta_{\lambda^*}^{-1})c_{\lambda^*,\lambda} = \varphi S_B j_\lambda. \end{aligned}$$

Therefore, since $\mathrm{id}_B = \sum_{\lambda\in\Lambda} j_\lambda p_\lambda$, we get that $\varepsilon_{\pi(A)}\varphi = \varepsilon_B$, $\Delta_{\pi(A)}\varphi = (\varphi\otimes\varphi)\Delta_B$, $m_{\pi(A)}(\varphi\otimes\varphi) = \varphi m_B$, and $S_{\pi(A)}\varphi = \varphi S_B$. Finally, we conclude by remarking that $\varphi\eta_B = \varphi j_{\mathbb{1}} = \pi(i_{\mathbb{1}}) = \pi(\eta_A) = \eta_{\pi(A)}$. □

Corrolary 3.9. *Let $\beta \in \mathrm{Hom}_{\mathcal{B}}(\mathbb{1}, B)$ and $\alpha \in \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, A)$ such that $\pi(\alpha) = \varphi\beta$.*

- (a) *If $\beta \in \mathcal{I}(\mathcal{B})$, then $\alpha \in \mathcal{I}(\mathcal{C})$ and $\tau_{\mathcal{C}}(L;\alpha) = \tau_{\mathcal{B}}(L;\beta)$ for any framed link L.*
- (b) *If $\beta \in \mathcal{I}(\mathcal{B})^{\mathrm{norm}}$, then $\alpha \in \mathcal{I}(\mathcal{C})^{\mathrm{norm}}$ and $\tau_{\mathcal{C}}(M;\alpha) = \tau_{\mathcal{B}}(M;\beta)$ for any 3-manifold M.*

Proof. This is an immediate consequence of Lemma 3.8 and Proposition 2.7. □

Remark 3.10. By Corollary 3.9, we have that

$$\bigcup_{\mathcal{B}} \pi^{-1}\big(\varphi_{\mathcal{B}}(\mathcal{I}(\mathcal{B}))\big) \subset \mathcal{I}(\mathcal{C}) \quad \text{and} \quad \bigcup_{\mathcal{B}} \pi^{-1}\big(\varphi_{\mathcal{B}}(\mathcal{I}(\mathcal{B})^{\mathrm{norm}})\big) \subset \mathcal{I}(\mathcal{C})^{\mathrm{norm}},$$

where $\mathcal{B}$ runs over (equivalence classes of) finitely semisimple full ribbon and abelian subcategories of $\mathcal{C}^s$ whose simple objects are scalar, and $\varphi_{\mathcal{B}}$ is the morphism (3.10) corresponding to $\mathcal{B}$. We will see in Section 4 that these inclusions may be strict (see Remark 4.12). This means that the semisimplification process “lacks” some invariants.

4. The case of categories of representations

In this section, we focus on the case of the category rep_H of representations of a finite-dimensional ribbon Hopf algebra H. In particular, we describe $\mathcal{I}(\mathrm{rep}_H)$ in purely algebraic terms. One of the interest of such a description is to avoid the representation theory of H, which may be of wild type. Moreover, we show that the 3-manifolds invariants obtained with these Kirby elements can be computed by using the Kauffman-Radford algorithm (even in the non-unimodular case).

4.1. Finite-dimensional Hopf algebras.

All considered algebras are supposed to be over the field $\Bbbk$.

Let H be a finite-dimensional Hopf algebra. Recall that a *left* (resp. *right*) *integral* for H is an element $\Lambda \in H$ such that $x\Lambda = \varepsilon(x)\Lambda$ (resp. $\Lambda x = \varepsilon(x)\Lambda$) for all $x \in H$. A left (resp. right) integral for H^* is then an element $\lambda \in H^*$ such that $x_{(1)}\lambda(x_{(2)}) = \lambda(x)\,1$ (resp. $\lambda(x_{(1)})x_{(2)} = \lambda(x)\,1$) for all $x \in H$. Since H is finite-dimensional, the space of left (resp. right) integrals for H is one-dimensional, and there always exist non-zero right integral λ for H^* and a non-zero left integral Λ for H such that $\lambda(\Lambda) = \lambda(S(\Lambda)) = 1$, see [Rad90, Proposition 3].

By [Rad94b, Corollary 2], the space H^* endowed with the right H-action defined, for any $f \in H^*$ and $h, x \in H$, by

$$\langle f\cdot h, x\rangle = \langle f, hx\rangle, \tag{4.1}$$

is a free H-module of rank 1 with basis every non-zero right integral λ for H^*. Likewise, H endowed with the right H^*-action $\leftharpoonup$ defined, for any $f \in H^*$ and $x \in H$, by

$$x \leftharpoonup f = f(x_{(1)})x_{(2)}, \tag{4.2}$$

is a free H^*-module of rank 1 with basis $S(\Lambda)$, where Λ is a non-zero left integral for H.

Lemma 4.1. *Let λ be a right integral for H^* and Λ be a left integral for H such that $\lambda(\Lambda) = \lambda(S(\Lambda)) = 1$. Let $a \in H$ and $f \in H^*$. Then*

(a) $a = \lambda(a\Lambda_{(1)})\, S(\Lambda_{(2)}) = \lambda(S(\Lambda_{(2)})a)\, \Lambda_{(1)}$.
(b) $f = \lambda \cdot a$ *if and only if* $a = (f \otimes S)\Delta(\Lambda)$.

Proof. Let us prove Part (a). Since λ is a right integral for H^* and λ is a left integral for H such that $\lambda(\Lambda) = 1$, we have that

$$\begin{aligned}
\lambda(a\Lambda_{(1)})\, S(\Lambda_{(2)}) &= \lambda(a_{(1)}\Lambda_{(1)})\, a_{(2)}\Lambda_{(2)}\, S(\Lambda_{(3)}) \\
&= \lambda(a_{(1)}\Lambda_{(1)})\, a_{(2)}\varepsilon(\Lambda_{(2)}) \\
&= \lambda(a_{(1)}\Lambda)\, a_{(2)} \\
&= \lambda(\Lambda)\, \varepsilon(a_{(1)})a_{(2)} \\
&= a.
\end{aligned}$$

The second equality of Part (a) can be proved similarly (by using the facts that λ is a right integral for H^* and $S(\Lambda)$ is a right integral for H such that $\lambda(S(\Lambda)) = 1$).

Let us prove Part (b). If $f = \lambda \cdot a$ then, by Part (a),

$$a = \lambda(a\Lambda_{(1)})\, S(\Lambda_{(2)}) = f(\Lambda_{(1)})\, S(\Lambda_{(2)}) = (f \otimes S)\Delta(\Lambda).$$

Conversely, if $a = (f \otimes S)\Delta(\Lambda)$ then, by Part (a),

$$\lambda(ax) = \lambda(f(\Lambda_{(1)})\, S(\Lambda_{(2)})x) = f(\lambda(S(\Lambda_{(2)})x)\Lambda_{(1)}) = f(x)$$

for all $x \in H$, and so $f = \lambda \cdot a$. □

Recall that an element $h \in H$ is *grouplike* if $\Delta(g) = g \otimes g$ and $\varepsilon(g) = 1$. We denote by $G(H)$ the space of grouplike elements of H. Recall (see [Abe80]) that there exist a unique grouplike element g of H such that $x_{(1)}\lambda(x_{(2)}) = \lambda(x)\, g$ for any $x \in H$ and any right integral $\lambda \in H^*$, and a unique grouplike element $\nu \in G(H^*) = \mathrm{Alg}(H, \Bbbk)$ such that $\Lambda x = \nu(x)\Lambda$ for any $x \in H$ and any left integral $\Lambda \in H$. The element $g \in H$ (resp. $\nu \in H^*$) is called the *distinguished grouplike element* of H (resp. of H^*). The Hopf algebra H is said to be *unimodular* if its integrals are to sided, that is, if $\nu = \varepsilon$.

By [Rad94b, Theorem 3] and [Rad94b, Proposition 3], right integrals for H^* and distinguished grouplike elements of H and H^* are related by:

$$\lambda(xy) = \lambda(S^2(y \leftharpoonup \nu)x) = \lambda((S^2(y) \leftharpoonup \nu)x) \quad \text{for all } x, y \in H, \tag{4.3}$$

$$\lambda(S(x)) = \lambda(gx) \quad \text{for all } x \in H. \tag{4.4}$$

4.2. **Quasitriangular Hopf algebras.** Following [Dri90], a Hopf algebra H is *quasitriangular* if it is endowed with an invertible element $R \in H \otimes H$ (the *R-matrix*) such that, for any $x \in H$,

$$R\Delta(x) = \sigma\Delta(x)R, \tag{4.5}$$

$$(\mathrm{id}_H \otimes \Delta)(R) = R_{13}R_{12}, \tag{4.6}$$

$$(\Delta \otimes \mathrm{id}_H)(R) = R_{13}R_{23}. \tag{4.7}$$

where $\sigma : H \otimes H \to H \otimes H$ denotes the usual flip map. The R-matrix verifies:

$$(\varepsilon \otimes \mathrm{id}_H)(R) = (\mathrm{id}_H \otimes \varepsilon)(R) = 1, \tag{4.8}$$

$$(S \otimes \mathrm{id}_H)(R) = R^{-1} = (\mathrm{id}_H \otimes S^{-1})(R), \tag{4.9}$$

$$R_{23}R_{13}R_{12} = R_{12}R_{13}R_{23}. \tag{4.10}$$

The *Drinfeld element* u associated to R is $u = m(S \otimes \mathrm{id}_H)\sigma(R) \in H$. It is invertible and verifies:

$$u^{-1} = m(\mathrm{id}_H \otimes S^2)(R_{21}), \tag{4.11}$$

$$S^2(x) = uxu^{-1} \quad \text{for all } x \in H, \tag{4.12}$$

$$\Delta(u) = (R_{21}R)^{-1}(u \otimes u), \tag{4.13}$$

$$\varepsilon(u) = 1. \tag{4.14}$$

Let H be a finite-dimensional quasitriangular Hopf algebra and let $\nu \in H^*$ be the distinguished grouplike element of H^*. Set $h_\nu = (\mathrm{id}_H \otimes \nu)(R) \in H$. By (4.7) and (4.8), h_ν is grouplike.

4.3. **Ribbon Hopf algebras.** Following [RT91], a *ribbon* Hopf algebra is a quasitriangular Hopf algebra H endowed with an invertible element $\theta \in H$ (the *twist*) such that:

$$\theta \text{ is central}, \tag{4.15}$$

$$S(\theta) = \theta, \tag{4.16}$$

$$\Delta(\theta) = (\theta \otimes \theta)R_{21}R. \tag{4.17}$$

The twist verifies:

$$\varepsilon(\theta_1) = 1, \tag{4.18}$$

$$\theta^{-2} = uS(u) = S(u)u. \tag{4.19}$$

Set $G = \theta u \in H$. Then G is grouplike and verifies:

$$S(u) = G^{-1}uG^{-1}, \tag{4.20}$$

$$S^2(x) = GxG^{-1}. \tag{4.21}$$

The element G is called the *special grouplike element* of H.

By [Rad92, Theorem 2] and (4.20), the special grouplike element G of a finite-dimensional ribbon Hopf algebra H is related to the distinguished grouplike element g of H and to $h_\nu = (\mathrm{id}_H \otimes \nu)(R) \in G(H)$ by

$$g = G^2 h_\nu. \tag{4.22}$$

Relations (4.4) and (4.22) imply that

$$\lambda(S(x)) = \lambda(G^2 h_\nu x) \quad \text{for all } x \in H. \tag{4.23}$$

4.4. **Category of representations of a ribbon Hopf algebra.** Let H be a ribbon Hopf algebra with R-matrix R and twist θ. Denote by rep_H the $\Bbbk$-category of finite-dimensional left H-modules and H-linear homomorphisms. The category rep_H is a monoidal $\Bbbk$-category with tensor product and unit object defined in the usual way using the comultiplication and counit of H. The category rep_H posses a left duality: for any module $M \in \mathrm{rep}_H$, set $M^* = \mathrm{Hom}_\Bbbk(M, \Bbbk)$, where $h \in H$ acts as the transpose of $x \in M \mapsto S(h)x \in M$. The duality morphism $\mathrm{ev}_M : M^* \otimes M \to \mathbb{1} = \Bbbk$ is the evaluation pairing and, if $(e_k)_k$ is a basis of M with dual basis $(e_k^*)_k$, then $\mathrm{coev}(1_\Bbbk) = \sum_k e_k \otimes e_k^*$. The category rep_H is braided: for modules $M, N \in \mathrm{rep}_H$, the braiding $c_{M,N} : M \otimes N \to N \otimes M$ is the composition of multiplication by R and the flip map $\sigma_{M,N} : M \otimes N \to N \otimes M$. The category rep_H is ribbon: for any module $M \in \mathrm{rep}_H$, the twist $\theta_M : M \to M$ is the multiplication by θ. Recall, see (1.9) and (1.10), that rep_H posses a right duality $M \in \mathrm{rep}_H \mapsto (M^*, \widetilde{\mathrm{ev}}_M, \widetilde{\mathrm{coev}}_M)$. Finally, the $\Bbbk$-category rep_H is pure, that is, $\mathrm{End}_{\mathrm{rep}_H}(\Bbbk) = \Bbbk$. Hence rep_H is a ribbon $\Bbbk$-category in the sense of Section 1.2.

Lemma 4.2. *Let G be the special grouplike element of H and M be a finite-dimensional left H-module. Then,*

(a) $\widetilde{\mathrm{ev}}_M(m \otimes f) = f(Gm)$ *for any* $f \in M^*$ *and* $m \in M$.
(b) $\widetilde{\mathrm{coev}}_M = (\mathrm{id}_{M^*} \otimes G^{-1}\mathrm{id}_M)\sigma_{M,M^*}\mathrm{coev}_M$.

Proof. Write $R = \sum_i a_i \otimes b_i$. Recall that $u = \sum_i S(b_i)a_i$. Then

$$\begin{aligned}\widetilde{\mathrm{ev}}_M(m \otimes f) &= \mathrm{ev}_M\, c_{M,M^*}(\theta_M \otimes \mathrm{id}_{M^*})(m \otimes f)\\ &= \textstyle\sum_i \mathrm{ev}_M(b_i f \otimes a_i\theta m) = \sum_i f(S(b_i)a_i\theta m)\\ &= f(u\theta m) = f(Gm).\end{aligned}$$

Let $(e_k)_k$ be a basis of M and $(e_k^*)_k$ be its dual basis. Note that if g is any $\Bbbk$-linear endomorphism of M, then $\sum_k g^*(e_k^*) \otimes e_k = \sum_k e_k^* \otimes g(e_k)$. For any $h \in H$, denote by $\rho(h)$ the $\Bbbk$-linear endomorphism of M defined by $m \in M \mapsto hm \in M$. By (4.9) and (4.11), we have that $R^{-1} = \sum_i S(a_i) \otimes b_i$ and $u^{-1} = \sum_i b_i S^2(a_i)$. Then

$$\begin{aligned}\widetilde{\mathrm{coev}}_M(1_\Bbbk) &= (\mathrm{id}_{M^*} \otimes \theta_M^{-1})(c_{M^*,M})^{-1}\mathrm{coev}_M(1_\Bbbk)\\ &= \textstyle\sum_{k,i} S(a_i)e_k^* \otimes \theta^{-1}b_i \cdot e_k\\ &= \textstyle\sum_i \big(\mathrm{id}_{M^*} \otimes \rho(\theta^{-1}b_i)\big)\big(\sum_k \rho(S^2(a_i))^*(e_k^*) \otimes e_k\big)\\ &= \textstyle\sum_i \big(\mathrm{id}_{M^*} \otimes \rho(\theta^{-1}b_i)\big)\big(\sum_k e_k^* \otimes \rho(S^2(a_i))(e_k)\big)\\ &= \big(\mathrm{id}_{M^*} \otimes \rho(\theta^{-1}\textstyle\sum_i b_i S^2(a_i))\big)\big(\sum_k e_k^* \otimes e_k\big)\\ &= \big(\mathrm{id}_{M^*} \otimes \rho(\theta^{-1}u^{-1})\big)\sigma_{M,M^*}\mathrm{coev}_M(1_k)\\ &= \big(\mathrm{id}_{M^*} \otimes \rho(G^{-1})\big)\sigma_{M,M^*}\mathrm{coev}_M(1_k)\\ &= (\mathrm{id}_{M^*} \otimes G^{-1}\mathrm{id}_M)\sigma_{M,M^*}\mathrm{coev}_M(1_k).\end{aligned}$$

This completes the proof of the lemma. □

We immediately deduce from Lemma 4.2 that, in the category rep_H, we have

$$\mathrm{tr}_q(f) = \mathrm{Tr}(Gf) = \mathrm{Tr}(G^{-1}f),$$
$$\dim_q(M) = \mathrm{Tr}(G\,\mathrm{id}_M) = \mathrm{Tr}(G^{-1}\mathrm{id}_M),$$

for all module $M \in \mathrm{rep}_H$ and all H-linear endomorphism f of M, where Tr denotes the usual trace of $\Bbbk$-linear endomorphisms.

4.5. **Braided Hopf algebra associated to ribbon Hopf algebras.** Let H be a finite-dimensional ribbon Hopf algebra. The ribbon $\Bbbk$-category rep_H of finite-dimensional left H-modules posses a coend (A, i) for the functor (1.16). More precisely, $A = H^* = \mathrm{Hom}_\Bbbk(H, \Bbbk)$ as a $\Bbbk$-space and is endowed with the coadjoint left H-action $\triangleright$ given, for any $f \in A$ and $h, x \in H$, by

$$\langle h \triangleright f, x\rangle = \langle f, S(h_{(1)})xh_{(2)}\rangle, \tag{4.24}$$

where $\langle , \rangle$ denotes the usual pairing between a $\Bbbk$-space and its dual. Given a module $M \in \mathrm{rep}_H$, the map $i_M : M^* \otimes M \to A$ is given by

$$\langle i_M(l \otimes m), x\rangle = \langle l, xm\rangle, \tag{4.25}$$

for all $l \in M^*$, $m \in M$, and $x \in H$.

Lemma 4.3. *If ξ is a dinatural transformation from the functor* (1.16) *of* rep_H *to a module $Z \in \mathrm{rep}_H$, then the (unique) morphism $r : A \to Z$ such that $\xi_M = r \circ i_M$ for all $M \in \mathrm{rep}_H$ is given by $f \in A = H^* \mapsto r(f) = \xi_H(f \otimes 1_H)$.*

Recall (see Section 1.6) that A is a Hopf algebra in rep_H. Using Lemma 4.3, the structural morphisms of A can be explicitly described in terms of the structure maps of the Hopf algebra H. Nevertheless, it is more convenient to write down its pre-dual structural morphisms: for example, since $A = H^*$ as a $\Bbbk$-space and H is finite-dimensional, the pre-dual of the multiplication $m_A : A \otimes A \to A$ of A is a morphism $\Delta^{\mathrm{Bd}} : H \to H \otimes H$ such that $(\Delta^{\mathrm{Bd}})^* = m_A$. That yields a $\Bbbk$-space $H^{\mathrm{Bd}} = H$ endowed with a comultiplication $\Delta^{\mathrm{Bd}} : H^{\mathrm{Bd}} \to H^{\mathrm{Bd}} \otimes H^{\mathrm{Bd}}$, a counit $\varepsilon^{\mathrm{Bd}} : H_1^{\mathrm{Bd}} \to \Bbbk$, a unit $\eta^{\mathrm{Bd}} : \Bbbk \to H^{\mathrm{Bd}}$, a multiplication $m^{\mathrm{Bd}} : H^{\mathrm{Bd}} \otimes H^{\mathrm{Bd}} \to H^{\mathrm{Bd}}$, and an antipode $S^{\mathrm{Bd}} : H^{\mathrm{Bd}} \to H^{\mathrm{Bd}}$. These structure maps, described in the following lemma, verify the same axioms as those of a Hopf algebra except that the usual flip maps are replaced by the braiding of rep_H. The space H^{Bd} is called the *braided Hopf algebra associated to H*, see [Maj93, Lyu95b].

Lemma 4.4 (cf. [Lyu95a])**.** *The braided Hopf algebra H^{Bd} associated to H can be described as follows :*

- *$H^{\mathrm{Bd}} = H$ as an algebra;*
- $\Delta^{\mathrm{Bd}}(x) = \sum_i x_{(2)}a_i \otimes S(b_{i(1)})x_{(1)}b_{i(2)} = \sum_i S(a_{i(1)})x_{(1)}a_{i(2)} \otimes S(b_i)x_{(2)}$,
- *$\varepsilon^{\mathrm{Bd}} = \varepsilon$;*
- $S_\alpha^{\mathrm{Bd}}(x) = \sum_i S(a_i)\theta^2 S(x)ub_i = \sum_i S(a_i)S(x)S(u)^{-1}b_i$,

for any $x \in H$, where $R = \sum_i a_i \otimes b_i$ is the R-matrix, u the Drinfeld element, and θ the twist of H.

4.6. **Kirby elements of ribbon Hopf algebras.** Throughout this section, H will denote a finite-dimensional ribbon Hopf algebra with R-matrix $R \in H \otimes H$ and twist $\theta \in H$. Let $u \in H$ be the Drinfeld element of H, $G = \theta u$ be the special grouplike element of H, $\lambda \in H^*$ be a non-zero right integral for H^*, $\Lambda \in H$ be a non-zero left integral for H such that $\lambda(\Lambda) = \lambda(S(\Lambda)) = 1$, $g \in G(H)$ be the distinguished grouplike element of H, $\nu \in G(H^*) = \mathrm{Alg}(H, \Bbbk)$ be the distinguished grouplike element of H^*, and $h_\nu = (\mathrm{id}_H \otimes \nu)(R) \in G(H)$. Let (A, i) be the coend for the functor (1.16) of rep_H, and let H^{Bd} be the braided Hopf algebra associated to H. Recall that $A = H^*$ endowed with the coadjoint left H-action $\triangleright$ and that A is a Hopf algebra in the category rep_H whose structure maps are dual to those of H^{Bd}. Recall that $\cdot$ denotes the right action of H on H^* defined in (4.1) and $\leftharpoonup$

denotes the right H^*-action on H defined in (4.2).

Set:

$$\phi : \begin{cases} H & \to \mathrm{Hom}_{\Bbbk}(\Bbbk, H^*) \\ z & \mapsto \phi_z = (\alpha \mapsto \alpha\lambda \cdot z) \end{cases}, \qquad T : \begin{cases} H & \to H \\ z & \mapsto T(z) = (S(z) \leftharpoonup \nu)h_\nu \end{cases},$$

and

$$\psi : \begin{cases} H^{\otimes n} & \to \mathrm{Hom}_{\Bbbk}(H^{*\otimes n}, \Bbbk) \\ X & \mapsto \psi_X = \big(f_1 \otimes \cdots \otimes f_n \mapsto \langle f_1 \otimes \cdots \otimes f_n, X\rangle\big) \end{cases}.$$

Denote by $\triangleleft$ be the right action of H on $H^{\otimes n}$ given by

$$(x_1 \otimes \cdots \otimes x_n) \triangleleft h = S(h_{(1)})x_1h_{(2)} \otimes \cdots \otimes S(h_{(2n-1)})x_nh_{(2n)}$$

for any $h \in H$ and $x_1, \ldots, x_n \in H$. Set

$$\begin{aligned} L(H) &= \{z \in H \mid (x \leftharpoonup \nu)z = zx \text{ for any } x \in H\}, \\ N(H) &= \{z \in L(H) \mid \lambda(za) = 0 \text{ for any } a \in Z(H)\}, \\ V_n(H) &= \{X \in H^{\otimes n} \mid X \triangleleft h = \varepsilon(h)X \text{ for any } h \in H\}. \end{aligned}$$

Note that if H is unimodular, then $L(H) = Z(H)$ and $T(z) = S(z)$ for all $z \in H$. By the definition of $L(H)$ and by (4.3), we have that

$$\lambda(zxy) = \lambda(zS^2(y)x) \tag{4.26}$$

for all $z \in L(H)$ and $x, y \in H$.

Lemma 4.5. (a) *The map ϕ is an $\Bbbk$-isomorphism with inverse given by:*

$$\alpha \in \mathrm{Hom}_{\Bbbk}(\Bbbk, H^*) \mapsto \phi^{-1}(\alpha) = (\alpha(1_{\Bbbk}) \otimes S)\Delta(\Lambda) \in H.$$

(b) *The map ϕ induces a $\Bbbk$-isomorphism between $L(H)$ and $\mathrm{Hom}_{\mathrm{rep}_H}(\Bbbk, A)$.*
(c) *The map ϕ induces a $\Bbbk$-isomorphism between $N(H)$ and $\mathrm{Negl}_{\mathrm{rep}_H}(\Bbbk, A)$.*
(d) *The map ψ induces a $\Bbbk$-isomorphism between $V_n(H)$ and $\mathrm{Hom}_{\mathrm{rep}_H}(A^{\otimes n}, \Bbbk)$.*
(e) *$V_1(H) = Z(H)$.*

Remark 4.6. In Appendix A, we use the space $L(H)$ and the morphism T, defined from topological considerations, to parameterize all the traces of a finite-dimensional ribbon Hopf algebra H.

Proof. Let us prove Part (a). Since $(H^*, \cdot)$ is a free right H-module of rank 1 with basis λ, ϕ is an isomorphism. The expression of ϕ^{-1} follows from Lemma 4.1(b).

Let us prove Part (b). Let $z \in H$. We have to show that $z \in L(H)$ if and only if ϕ_z is H-linear. Suppose that ϕ_z is H-linear. For all $a, h \in H$,

$$\begin{aligned} \varepsilon(h)\,\lambda(za) &= \langle \varepsilon(S^{-1}(h))\,\phi_z(1_{\Bbbk}), a\rangle = \langle S^{-1}(h) \triangleright \phi_z(1_{\Bbbk}), a\rangle \\ &= \lambda(zh_{(2)}aS^{-1}(h_{(1)})) = \lambda(S^2(S^{-1}(h_{(1)}) \leftharpoonup \nu)zh_{(2)}a) \quad \text{by (4.3)}, \end{aligned}$$

and so, since $(H^*, \cdot)$ is a free right H-module of rank 1 with basis λ,

$$\varepsilon(h)z = S^2(S^{-1}(h_{(1)}) \leftharpoonup \nu)zh_{(2)} = \nu^{-1}(h_{(2)})\,S(h_{(1)})zh_{(3)} \tag{4.27}$$

for all $h \in H$. Hence, for any $x \in H$,

$$\begin{aligned}(x \leftharpoonup \nu)z &= \nu(x_{(1)})\, x_{(2)} z = \nu(x_{(1)})\, x_{(2)}\varepsilon(x_{(3)}) z \\ &= \nu(x_{(1)})\, \nu^{-1}(x_{(4)})\, x_{(2)} S(x_{(3)}) z x_{(5)} \quad \text{by (4.27)} \\ &= \nu(x_{(1)})\, \nu^{-1}(x_{(3)})\, \varepsilon(x_{(2)})\, z x_{(4)} = \nu(x_{(1)})\, \nu^{-1}(x_{(2)})\, z x_{(3)} \\ &= \varepsilon(x_{(1)})\, z x_{(2)} = zx,\end{aligned}$$

and so $z \in L(H)$. Conversely, suppose that $z \in L(H)$. Then, for any $x, h \in H$,

$$\begin{aligned}\lambda(zS(h_{(1)})xh_{(2)}) &= \lambda(zS^2(h_{(2)})S(h_{(1)})x) \quad \text{by (4.26)} \\ &= \lambda(zS(h_{(1)}S(h_{(2)}))x) = \varepsilon(h)\, \lambda(zx),\end{aligned}$$

and so ϕ_z is H-linear.

Let us prove Part (d). Note that ψ is an isomorphism since H is finite-dimensional. Let $X \in H^{\otimes n}$. For all $h \in H$ and $f_1, \dots, f_n \in H^*$,

$$\psi_X\big(h \triangleright (f_1 \otimes \cdots \otimes f_n)\big) = \langle f_1 \otimes \cdots \otimes f_n, X \triangleleft h\rangle$$

and

$$\varepsilon(h)\, \psi_X(f_1 \otimes \cdots \otimes f_n) = \langle f_1 \otimes \cdots \otimes f_n, \varepsilon(h)\, X\rangle.$$

Therefore ψ_X is H-linear if and only if $X \in V_n(H)$.

Let us prove Part (e). Let $a \in Z(H)$. For all $h \in H$,

$$a \triangleleft h = S(h_{(1)})ah_{(2)} = S(h_{(1)})h_{(2)}a = \varepsilon(h)\, a$$

and so $a \in V_1(H)$. Conversely, let $a \in V_1(H)$. For all $x \in H$,

$$xa = x_{(1)}\varepsilon(x_{(2)})\, a = x_{(1)}(a \triangleleft x_{(2)}) = x_{(1)}S(x_{(2)})ax_{(3)} = \varepsilon(x_{(1)})\, ax_{(2)} = ax,$$

and so $a \in Z(H)$.

Finally, let us prove Part (c). Let $z \in L(H)$. Since $\mathrm{End}_{\mathrm{rep}_H}(\Bbbk) = \Bbbk$, we have that ϕ_z is negligible if and only if $\psi_a\phi_z = \lambda(za) = 0$ for all $a \in V_1(H) = Z(H)$, that is, if and only if $z \in N(H)$. □

Lemma 4.7. (a) *$L(H)$ is a commutative algebra with product $*$ defined by*

$$\begin{aligned}x * z &= \lambda(xS(z_{(2)}))\, z_{(1)} = \lambda(zS(x_{(2)}))\, x_{(1)} \\ &= \lambda(z_{(1)}S^{-1}(x))\, z_{(2)} = \lambda(x_{(1)}S^{-1}(z))\, x_{(2)}\end{aligned}$$

for any $x, z \in L(H)$, and with $S(\Lambda)$ as unit element.

(b) *For any $z \in L(H)$, $S_A \circ \phi_z = \phi_{T(z)}$, where S_A denotes the antipode of the categorical Hopf algebra A.*

(c) *T induces on $L(H)$ an involutory algebra-automorphism, that is,*

$$T(x * z) = T(x) * T(z), \quad T(S(\Lambda)) = S(\Lambda), \quad \textit{and} \quad T^2(x) = x$$

for all $x, z \in L(H)$.

Proof. Let us prove Part (a). Since A is a Hopf algebra in rep_H, the space $\mathrm{Hom}_{\mathrm{rep}_H}(\Bbbk, A)$ is an algebra for the convolution product $\alpha * \beta = m_A(\alpha \otimes \beta)$ and with unit element η_A. This algebra structure transports to $L(H)$ via the $\Bbbk$-isomorphism $\phi : L(H) \to \mathrm{Hom}_{\mathrm{rep}_H}(\Bbbk, A)$. Let $x, z \in L(H)$. Then

$$\begin{aligned}x * z &= \phi^{-1}(\phi_x * \phi_z) = \phi^{-1}(m_A(\phi_x \otimes \phi_z)) \\ &= \langle m_A(\phi_x \otimes \phi_z)(1_\Bbbk), \Lambda_{(1)}\rangle\, S(\Lambda_{(2)}) \quad \text{by Lemma 4.5(a)} \\ &= \langle \lambda \cdot x \otimes \lambda \cdot z, \Delta^{\mathrm{Bd}}(\Lambda_{(1)})\rangle\, S(\Lambda_{(2)})\end{aligned}$$

Write $R=\sum_i a_i\otimes b_i$. By using Lemma 4.4, (4.8) and (4.26), we have that

$$\begin{aligned} x*z &= \textstyle\sum_i \lambda(x\Lambda_{(2)}a_i)\,\lambda(zS(b_{i(1)})\Lambda_{(1)}b_{i(2)})\,S(\Lambda_{(3)})\\ &= \textstyle\sum_i \lambda(x\Lambda_{(2)}a_i)\,\lambda(zS^2(b_{i(2)})S(b_{i(1)})\Lambda_{(1)})\,S(\Lambda_{(3)})\\ &= \textstyle\sum_i \lambda(x\Lambda_{(2)}a_i\varepsilon(b_i))\,\lambda(z\Lambda_{(1)})\,S(\Lambda_{(3)}), \end{aligned}$$

that is,

$$x*z=\lambda(x\Lambda_{(2)})\,\lambda(z\Lambda_{(1)})\,S(\Lambda_{(3)}). \tag{4.28}$$

Likewise

$$\begin{aligned} x*z &= \textstyle\sum_i \lambda(xS(a_{i(1)})\Lambda_{(1)}a_{i(2)})\,\lambda(zS(b_i)\Lambda_{(2)})\,S(\Lambda_{(3)})\\ &= \textstyle\sum_i \lambda(xS^2(a_{i(2)})S(a_{i(1)})\Lambda_{(1)})\,\lambda(zS(b_i)\Lambda_{(2)})\,S(\Lambda_{(3)})\\ &= \textstyle\sum_i \lambda(x\Lambda_{(1)})\,\lambda(zS(\varepsilon(a_i)\,b_i)\Lambda_{(2)})\,S(\Lambda_{(3)}), \end{aligned}$$

that is,

$$x*z=\lambda(x\Lambda_{(1)})\,\lambda(z\Lambda_{(2)})\,S(\Lambda_{(3)}). \tag{4.29}$$

Now, by Lemma 4.1(a),

$$z=\lambda(z\Lambda_{(1)})\,S(\Lambda_{(2)}), \tag{4.30}$$

$$x=\lambda(x\Lambda_{(1)})\,S(\Lambda_{(2)}), \tag{4.31}$$

so that

$$z_{(1)}\otimes S^{-1}(z_{(2)})=\lambda(z\Lambda_{(1)})\,S(\Lambda_{(3)})\otimes\Lambda_{(2)} \tag{4.32}$$

$$x_{(1)}\otimes S^{-1}(x_{(2)})=\lambda(x\Lambda_{(1)})\,S(\Lambda_{(3)})\otimes\Lambda_{(2)}. \tag{4.33}$$

Hence

$$\begin{aligned} x*z &= \lambda(x\Lambda_{(2)})\,\lambda(z\Lambda_{(1)})\,S(\Lambda_{(3)}) \quad \text{by (4.28)}\\ &= \lambda(xS^{-1}(z_{(2)}))\,z_{(1)} \quad \text{by (4.32)}\\ &= \lambda(xS(z_{(2)}))\,z_{(1)} \quad \text{by (4.26)}, \end{aligned}$$

$$\begin{aligned} x*z &= \lambda(z\Lambda_{(1)})\,\lambda(x\Lambda_{(2)})\,S(\Lambda_{(3)}) \quad \text{by (4.28)}\\ &= \lambda(z\Lambda_{(1)})\,\lambda(x_{(1)}\Lambda_{(2)})\,x_{(2)}\Lambda_{(3)}S(\Lambda_{(4)})\\ &= \lambda(x_{(1)}S^{-1}(z))\,x_{(2)} \quad \text{by (4.30)}, \end{aligned}$$

$$\begin{aligned} x*z &= \lambda(x\Lambda_{(1)})\,\lambda(z\Lambda_{(2)})\,S(\Lambda_{(3)}) \quad \text{by (4.29)}\\ &= \lambda(zS^{-1}(x_{(2)}))\,x_{(1)} \quad \text{by (4.33)}\\ &= \lambda(zS(x_{(2)}))\,x_{(1)} \quad \text{by (4.26)}, \end{aligned}$$

and

$$\begin{aligned} x*z &= \lambda(x\Lambda_{(1)})\,\lambda(z\Lambda_{(2)})\,S(\Lambda_{(3)}) \quad \text{by (4.29)}\\ &= \lambda(x\Lambda_{(1)})\,\lambda(z_{(1)}\Lambda_{(2)})\,z_{(2)}\Lambda_{(3)}S(\Lambda_{(4)})\\ &= \lambda(z_{(1)}S^{-1}(x))\,z_{(2)} \quad \text{by (4.31)}. \end{aligned}$$

Note that these expressions of the product of $L(H)$ show that $L(H)$ is commutative. Moreover, by Lemma 4.5(a), the unit element of $L(H)$ is

$$\phi^{-1}(\eta_A)=(\eta_A(1)\otimes S)\Delta(\Lambda)=(\varepsilon\otimes S)\Delta(\Lambda)=S(\Lambda).$$

Let us prove Part (b). Let $z \in L(H)$. Write $R = \sum_i a_i \otimes b_i$. For any $x \in H$,

$$\begin{aligned}
\langle S_A\phi_z(1_{\Bbbk}), x\rangle &= \langle \lambda \cdot z, S^{\mathrm{Bd}}(x)\rangle \\
&= \textstyle\sum_i \lambda(zS(a_i)\theta^2 S(x)ub_i) \quad \text{by Lemma 4.4} \\
&= \textstyle\sum_i \lambda(zS^2(u)S^2(b_i)S(a_i)\theta^2 S(x)) \quad \text{by (4.26)} \\
&= \lambda(zu^2\theta^2 S(x)) \quad \text{since } S^2(u) = u \text{ and } (S\otimes S)(R) = R \\
&= \lambda(zG^2 S(x)) = \lambda((G^2 \leftharpoonup \nu)zS(x)) = \nu(G^2)\,\lambda(G^2 zS(x)) \\
&= \nu(G^2)\,\lambda(G^2 h_\nu x S^{-1}(z)G^{-2}) \quad \text{by (4.23)} \\
&= \nu(G^2)\,\lambda((G^{-2} \leftharpoonup \nu)G^2 h_\nu x S^{-1}(x)) \quad \text{by (4.3)} \\
&= \nu(G^2)\,\nu(G^{-2})\,\lambda(h_\nu x S^{-1}(z)) \\
&= \lambda((S(z) \leftharpoonup \nu)h_\nu x) \quad \text{by (4.3)} \\
&= \lambda(T(z)x) = \langle \phi_{T(z)}(1_{\Bbbk}), x\rangle,
\end{aligned}$$

that is $S_A\phi_z = \phi_{T(z)}$.

Let us prove Part (c). Let $z \in L(H)$. Firstly $T(z) \in L(H)$ since $\phi_{T(z)} = S_A\phi_z$ is H-linear. Moreover, since $S_A^2 = \theta_A$ and the twist of rep_H is natural and verify $\theta_{\Bbbk} = \mathrm{id}_{\Bbbk}$, we have that

$$T^2(z) = \phi^{-1}(\phi_{T^2(z)}) = \phi^{-1}(S_A^2\phi_z) = \phi^{-1}(\theta_A\phi_z) = \phi^{-1}(\phi_z\theta_{\Bbbk}) = \phi^{-1}(\phi_z) = z.$$

For any $x, z \in L(H)$,

$$\begin{aligned}
\phi_{T(x*z)} &= S_A\phi_{x*z} = S_A(\phi_x * \phi_z) = S_A m_A(\phi_x \otimes \phi_z) \\
&= m_A(S_A \otimes S_A)c_{A,A}(\phi_x \otimes \phi_z) = m_A(S_A \otimes S_A)(\phi_z \otimes \phi_x)c_{\Bbbk,\Bbbk} \\
&= m_A(S_A\phi_z \otimes S_A\phi_x) = \phi_{T(z)} * \phi_{T(x)} = \phi_{T(z)*T(x)}
\end{aligned}$$

and so $T(x*z) = T(z)*T(x) = T(x)*T(z)$. Finally $T(S(\Lambda)) = S(\Lambda)$ since $\phi_{T(S(\Lambda))} = S_A\phi_{S(\Lambda)} = S_A\eta_A = \eta_A = \phi_{S(\Lambda)}$. □

In the next theorem, we describe the sets $\mathcal{I}(\mathrm{rep}_H)$ and $\mathcal{I}(\mathrm{rep}_H)^{\mathrm{norm}}$ in algebraic terms. Set

$$\mathcal{I}(H) = \phi^{-1}(\mathcal{I}(\mathrm{rep}_H)) \quad \text{and} \quad \mathcal{I}(H)^{\mathrm{norm}} = \phi^{-1}(\mathcal{I}(\mathrm{rep}_H)^{\mathrm{norm}}), \tag{4.34}$$

where $\phi : L(H) \to \mathrm{Hom}_{\mathrm{rep}_H}(k, A)$ is as in Lemma 4.5. Note that $\mathcal{I}(H)^{\mathrm{norm}} \subset \mathcal{I}(H)$. It follows from (2.3) that

$$\Bbbk\mathcal{I}(H) + N(H) \subset \mathcal{I}(H) \quad \text{and} \quad \Bbbk^*\mathcal{I}(H)^{\mathrm{norm}} + N(H) \subset \mathcal{I}(H)^{\mathrm{norm}}.$$

Since $\eta_A \in \mathcal{I}(\mathrm{rep}_H)^{\mathrm{norm}}$ and by Lemma 4.7, the right integral $S(\Lambda)$ for H belongs to $\mathcal{I}(H)$ and $\mathcal{I}(H)^{\mathrm{norm}}$.

Theorem 4.8. *$\mathcal{I}(H)$ is constituted by the element $z \in L(H)$ verifying*

(a) *$T(z) - z \in N(H)$, that is, $\lambda(T(z)a) = \lambda(za)$ for all $a \in Z(H)$;*

(b) *$\sum_i \lambda(zx_{i(1)})\lambda(zx_{i(2)}y_i) = \sum_i \lambda(zx_i)\lambda(zy_i)$ for all $X = \sum_i x_i \otimes y_i \in V_2(H)$.*

In particular, $\mathcal{I}(H)$ contains the elements $z \in L(H)$ satisfying $T(z) = z$ and $\lambda(zx_{(1)})zx_{(2)} = \lambda(zx)z$ for all $x \in H$. Moreover, an element $z \in \mathcal{I}(H)$ belongs to $\mathcal{I}(H)^{\mathrm{norm}}$ if and only if $\lambda(z\theta) \neq 0 \neq \lambda(z\theta^{-1})$.

Note that the sets $\mathcal{I}(H)$ and $\mathcal{I}(H)^{\mathrm{norm}}$ do not depend on the choice of the non-zero right integral λ for H^*. In Section 5, we give an example of determination of these sets for a family of non-unimodular ribbon Hopf algebras.

Proof. Let $z \in L(H)$. By Lemma 4.7(b), we have that $S_A\phi_z = \phi_{T(z)}$. Therefore, using Lemma 4.5(c), we get that $S_A\phi_z - \phi_z \in \mathrm{Negl}_{\mathrm{rep}_H}(\Bbbk, A)$ if and only if $T(z)-z \in N(H)$, that is, if and only if $\lambda(T(z)a) = \lambda(za)$ for all $a \in Z(H)$. Note that this last property is in particular satisfied when $T(z) = z$.

Since $\mathrm{End}_{\mathrm{rep}_H}(\Bbbk) = \Bbbk$, the morphism $\Gamma_r(\phi_z \otimes \phi_z) - \phi_z \otimes \phi_z : \Bbbk \to A \otimes A$ is negligible if and only if $G \circ \big(\Gamma_r(\phi_z \otimes \phi_z) - \phi_z \otimes \phi_z\big) = 0$ for any $G \in \mathrm{Hom}_{\mathrm{rep}_H}(A \otimes A, \Bbbk)$. By Lemma 4.5(d), this is equivalent to $\psi_X(\Gamma_r(\phi_z \otimes \phi_z) - \phi_z \otimes \phi_z) = 0$ for all $X \in V_2(H)$. Now, writing $R = \sum_i a_i \otimes b_i$ and using Lemma 4.4, we have that for any $x, y \in H$,

$$\begin{aligned}
&\langle \Gamma_r(\phi_z \otimes \phi_z)(1_\Bbbk), x \otimes y\rangle \\
&= \langle (m_A \otimes \mathrm{id}_A)(\mathrm{id}_A \otimes \Delta_A)(\lambda \cdot z \otimes \lambda \cdot z), x \otimes y\rangle \\
&= \langle (\mathrm{id}_A \otimes \Delta_A)(\lambda \cdot z \otimes \lambda \cdot z), \Delta^{\mathrm{Bd}}(x) \otimes y\rangle \\
&= \textstyle\sum_i \langle (\mathrm{id}_A \otimes \Delta_A)(\lambda \cdot z \otimes \lambda \cdot z), S(a_{i(1)})x_{(1)}a_{i(2)} \otimes S(b_i)x_{(2)} \otimes y\rangle \\
&= \textstyle\sum_i \langle \lambda \cdot z \otimes \lambda \cdot z, S(a_{i(1)})x_{(1)}a_{i(2)} \otimes S(b_i)x_{(2)}y\rangle \\
&= \textstyle\sum_i \lambda(zS(a_{i(1)})x_{(1)}a_{i(2)})\, \lambda(zS(b_i)x_{(2)}y) \\
&= \textstyle\sum_i \lambda(zS^2(a_{i(2)})S(a_{i(1)})x_{(1)})\, \lambda(zS(b_i)x_{(2)}y) \quad \text{by (4.26)} \\
&= \textstyle\sum_i \lambda(zx_{(1)})\, \lambda(zS(\varepsilon(a_i)b_i)x_{(2)}y) \\
&= \lambda(zx_{(1)})\, \lambda(zx_{(2)}y) \quad \text{by (4.8).}
\end{aligned}$$

Therefore the morphism $\Gamma_r(\phi_z \otimes \phi_z) - \phi_z \otimes \phi_z : \Bbbk \to A \otimes A$ is negligible if and only if $\lambda(zx_{i(1)})\, \lambda(zx_{i(2)}y_i) = \lambda(zx_i)\, \lambda(zy_i)$ for all $X = \sum x_i \otimes y_i \in V_2(H)$. Note that this last property is in particular satisfied when $\lambda(zx_{(1)})\, \lambda(zx_{(2)}y) = \lambda(zx)\, \lambda(zy)$ for all $x, y \in H$, that is, since $(H^*, \cdot)$ is a free right H-module of rank 1 with basis λ, when $\lambda(zx_{(1)})\, zx_{(2)} = \lambda(zx)\, z$ for all $x \in H$.

Finally, by using Lemma 4.3, we have that:

$$\Theta_\pm \phi_z = \mathrm{ev}_H(\mathrm{id}_{H^*} \otimes \theta_H^{\pm 1})(\phi_z(1_\Bbbk) \otimes 1_H) = \mathrm{ev}_H(\lambda \cdot z \otimes \theta^{\pm 1}) = \lambda(z\theta^{\pm 1}).$$

This completes the proof of the theorem. □

Corrolary 4.9. $1 \in \mathcal{I}(H)$ *if and only if* H *is unimodular.*

Proof. Suppose that H is unimodular. Therefore $L(H) = Z(H)$ and $T(z) = S(z)$ for all $z \in Z(H)$. In particular $1 \in L(H)$ and $T(1) = 1$. Moreover, $\lambda(x_{(1)})x_{(2)} = \lambda(x)1$ for all $x \in H$ (since λ is a right integral for H^*). Hence $1 \in \mathcal{I}(H)$ by Theorem 4.8.

Conversely, suppose that $1 \in \mathcal{I}(H)$. In particular $1 \in L(H)$ and so $z \leftharpoonup \nu = z$ for all $z \in H$. Therefore $\varepsilon(z) = \varepsilon(z \leftharpoonup \nu) = \nu(z_{(1)})\varepsilon(z_{(2)}) = \nu(z)$ for all $z \in H$, that is, H is unimodular. □

4.7. **Elements elements from semisimplification.** Let H be a finite-dimensional ribbon Hopf algebra. Let (A, i) be the coend of the functor (1.16) of rep_H (as in Section 4.5). Denote by rep_H^s the semisimplification of rep_H and by π its associated surjective ribbon functor $\mathrm{rep}_H \to \mathrm{rep}_H^s$ (see Section 3.4). Let $\phi : L(H) \to \mathrm{Hom}_{\mathrm{rep}_H}(\Bbbk, A)$ be as is Section 4.6. Set

$$\mathcal{I}(H)^s = \bigcup_{\mathcal{B}} \phi^{-1}\Big(\pi^{-1}\big(\varphi_{\mathcal{B}}(\mathcal{I}(\mathcal{B}))\big)\Big), \tag{4.35}$$

where $\mathcal{B}$ runs over (equivalence classes of) finitely semisimple full ribbon and abelian subcategories of rep^s_H whose simple objects are scalar, and $\varphi_{\mathcal{B}}$ is the morphism (3.10) corresponding to $\mathcal{B}$. By Corollary 3.9, we have that

$$\mathcal{I}(H)^s \subset \mathcal{I}(H).$$

Note that this inclusion may be strict (see Remark 4.12).

Let $\mathcal{V}$ be a set of representatives of isomorphism classes of indecomposable finite-dimensional left H-modules with non-zero quantum dimension. Note that $\pi(\mathcal{V}) = \{\pi(V) \,|\, V \in \mathcal{V}\}$ is a set of representatives of isomorphism classes of simple objects of rep^s_H. Let λ be a non-zero right integral for H^*. Since H^* is a free right H-module with basis λ (see Section 4.1), there exists a (unique) element $z_V \in H$ such that

$$\lambda(z_V x) = \mathrm{Tr}(G^{-1}x\,\mathrm{id}_V) \tag{4.36}$$

for all $x \in H$, where G is the special grouplike element of H. Recall that $\dim_q(V) = \mathrm{Tr}(G^{-1}\mathrm{id}_V)$ denotes the quantum dimension of V (see Lemma 4.2).

Corrolary 4.10. (a) *If $z \in \mathcal{I}(H)^s$, then $z = k\sum_{V\in\mathcal{W}} \dim_q(V)\, z_V$ for some finite subset $\mathcal{W}$ of $\mathcal{V}$ and some scalar $k \in \Bbbk$.*

(b) *Let $\mathcal{W}$ be a (finite) set of representatives of isomorphism classes of simple objects of a full ribbon and abelian subcategory of rep^s_H. We can suppose that $\mathcal{W} \subset \pi(\mathcal{V})$. If the objects of $\mathcal{W}$ are scalar, then*

$$\sum_{V\in\pi^{-1}(\mathcal{W})} \dim_q(V)\, z_V \in \mathcal{I}(H)^s.$$

Proof. Let $\mathcal{B}$ be finitely semisimple full ribbon and abelian subcategory of rep^s_H whose simple objects are scalar, and let (B, j) be the coend of the functor (1.16) of $\mathcal{B}$ (as in Section 3.2). We can suppose that there exists a (finite) subset $\mathcal{W}$ of $\mathcal{V}$ such that $\pi(\mathcal{W})$ is a set of representatives of isomorphism classes of simple objects of $\mathcal{B}$. Recall that $B = \oplus_{V\in\mathcal{W}}\, \pi(V)^* \otimes \pi(V)$. In particular, there exist morphisms $p_V : B \to \pi(V)^* \otimes \pi(V)$ and $q_V : \pi(V)^* \otimes \pi(V) \to B$ of $\mathcal{B}$ such that $\mathrm{id}_B = \sum_{V\in\mathcal{W}} q_V p_V$ and $p_V q_W = \delta_{V,W}\,\mathrm{id}_{\pi(V)^*\otimes\pi(V)}$. Recall that $j_V = q_V$ for any $V \in \mathcal{W}$. Let $\phi : L(H) \to \mathrm{Hom}_{\mathrm{rep}_H}(\Bbbk, A)$ be as is Section 4.6. As in (3.10), we set

$$\varphi_{\mathcal{B}} = \sum_{V\in\mathcal{W}} \pi(i_V)p_V \in \mathrm{Hom}_{\mathrm{rep}^s_H}(B, \pi(A)).$$

Let $V \in \mathcal{W}$. Let $(e_i)_i$ be a basis of V with dual basis $(e_i^*)_i$. By Lemma 4.2(b) and (4.25) we have that

$$\begin{aligned}
\langle i_V \widetilde{\mathrm{coev}}_V(1_\Bbbk), x\rangle &= \langle i_V(\mathrm{id}_{V^*} \otimes G^{-1}\mathrm{id}_V)\,\sigma_{V,V^*}\,\mathrm{coev}_V(1_\Bbbk), x\rangle\\
&= \textstyle\sum_i \langle i_V(e_i^* \otimes G^{-1}e_i), x\rangle = \sum_i \langle e_i^*, xG^{-1}e_i\rangle\\
&= \mathrm{Tr}(xG^{-1}\mathrm{id}_V) = \mathrm{Tr}(G^{-1}x\,\mathrm{id}_V)\\
&= \lambda(z_V x) = \langle \phi_{z_V}(1_\Bbbk), x\rangle,
\end{aligned}$$

for any $x \in H$, that is, $i_V\widetilde{\mathrm{coev}}_V = \phi_{z_V}$. Moreover,

$$\begin{aligned}
\varphi_{\mathcal{B}}\, j_{\pi(V)} \widetilde{\mathrm{coev}}_{\pi(V)} &= \sum_{W\in\mathcal{W}} \pi(i_W)p_W j_{\pi(V)} \widetilde{\mathrm{coev}}_{\pi(V)}\\
&= \sum_{W\in\mathcal{W}} \delta_{V,W}\, \pi(i_W)\pi(\widetilde{\mathrm{coev}}_V)\\
&= \pi(i_V\widetilde{\mathrm{coev}}_V) = \pi(\phi_{z_V}).
\end{aligned}$$

Hence Part (a) follows from Lemma 3.3 and Corollary 3.9(a), and Part (b) follows from Theorem 3.4 and Corollary 3.9(a). $\square$

Lemma 4.11. *If H is not semisimple, then $\varepsilon(z)=0$ for any $z \in \mathcal{I}(H)^s$.*

Remark 4.12. When H is not semisimple, it is possible that $\mathcal{I}(H)^s \subsetneq \mathcal{I}(H)$. For example, if H is unimodular but not semisimple, then $1 \in \mathcal{I}(H)$ (by Corollary 4.9) and $1 \notin \mathcal{I}(H)^s$ (by Lemma 4.11, since $\varepsilon(1)=1$).

Proof. Let Λ be a left integral for H such that $\lambda(\Lambda)=1$. Since H is not semisimple, we have $\varepsilon(\Lambda)=0$ (by [Abe80, Theorem 3.3.2]) and $\Lambda^2=\varepsilon(\Lambda)\Lambda=0$. Now, if M is a finite-dimensional left H-module, then $(\Lambda\,\mathrm{id}_M)^2=\Lambda^2\mathrm{id}_M=0$ and so $\mathrm{Tr}(\Lambda\,\mathrm{id}_M)=0$. Let $z \in \mathcal{I}(H)^s$. By Corollary 4.10(a), there exist $k \in \Bbbk$ and a finite subset $\mathcal{W}$ of $\mathcal{V}$ such that $z = k\sum_{V\in\mathcal{W}} \dim_q(V)\, z_V$. Then

$$\lambda(z\Lambda)=k\sum_{V\in\mathcal{W}}\dim_q(V)\,\mathrm{Tr}(G^{-1}\Lambda\,\mathrm{id}_V)=k\sum_{V\in\mathcal{W}}\dim_q(V)\varepsilon(G^{-1})\,\mathrm{Tr}(\Lambda\,\mathrm{id}_V)=0.$$

Hence $\varepsilon(z)=\varepsilon(z)\lambda(\Lambda)=\lambda(\varepsilon(z)\Lambda)=\lambda(z\Lambda)=0$. $\square$

Recall (see [CR62]) that $\Bbbk$ is a *splitting field* for a $\Bbbk$-algebra A if every simple finite dimensional left A-module is scalar. Note that this is always the case if $\Bbbk$ is algebraically closed.

Corrolary 4.13. *If H is semisimple and $\Bbbk$ is a splitting field for H, then $\mathcal{I}(H)^s = \mathcal{I}(H)$ and this set is composed by elements $z \in Z(H)$ satisfying $S(z)=z$ and $\lambda(zx_{(1)})zx_{(2)}=\lambda(zx)z$ for all $x \in H$.*

Proof. Since H is semisimple and finite-dimensional, we have that $\mathrm{rep}_H^s=\mathrm{rep}_H$ and that $\mathcal{V}$ finite. Moreover, since $\Bbbk$ is a splitting field for a H, every $V\in\mathcal{V}$ is scalar. Then $\mathcal{I}(H)=\phi^{-1}(\mathcal{I}(\mathrm{rep}_H))\subset\mathcal{I}(H)^s$ and so $\mathcal{I}(H)^s=\mathcal{I}(H)$. Moreover, since H is unimodular (because it is semisimple), we have that $L(H)=Z(H)$, $N(H)=0$, and $T(x)=S(x)$ for all $x\in H$ (see Section 4.6). Therefore, by using Theorem 4.8, we get that $z \in \mathcal{I}(H)$ if and only if $z\in Z(H)$, $S(z)=z$, and $\lambda(zx_{(1)})zx_{(2)}=\lambda(zx)z$ for all $x\in H$. $\square$

Proposition 4.14. *Suppose that H is semisimple, that $\Bbbk$ is of characteristic 0, and that $\Bbbk$ is a splitting field for H. Then $Z(H)=\bigoplus_{V\in\mathcal{V}}\Bbbk z_V$ and $\sum_{V\in\mathcal{V}}\dim_q(V)\,z_V \in \Bbbk^*1$.*

Proof. Note that H is cosemisimple since any finite-dimensional semisimple Hopf algebra over a field of characteristic 0 is cosemisimple (see [LR88, Theorem 3.3]). Then $S^2=\mathrm{id}_H$ by [LR87, Theorem 4] and $\lambda(1)\neq 0$ by [Abe80, Theorem 3.3.2].

Note that $\mathcal{V}$ is finite (since H is finite-dimensional). By [CR62, Theorem 25.10], any simple left ideal of H is isomorphic (as a left H-module) to a (unique) element of $\mathcal{V}$. For any $V\in\mathcal{V}$, let $H_V\subset H$ be the sum of all the simple left ideals of H which are isomorphic to V. By [CR62, Theorem 25.15], H_V is a two-sided ideal of H, H_V is simple $\Bbbk$-algebra (the operations being those induced by H), $B_VB_W=0$ for $V\neq W\in\mathcal{V}$, $H=\bigoplus_{V\in\mathcal{V}}H_V$, and H is isomorphic (as an algebra) to $\prod_{V\in\mathcal{V}}H_V$. Moreover, if e_V denotes the unit of H_V, then $1=\sum_{V\in\mathcal{V}}e_V$, $H_V=He_V$, and $e_Ve_W=\delta_{V,W}\,e_V$ for all $V,W\in\mathcal{V}$. By [CR62, Theorem 26.4], since H_V is a simple $\Bbbk$-algebra, V is a simple left H_V-module, and $\mathrm{End}_{H_V}(V)=\Bbbk$ (because V is a scalar H-module), we have that H_V is isomorphic (as an algebra) to $\mathrm{End}_\Bbbk(V)$ and that $\dim_\Bbbk(V)$ is the number of simple left ideals appearing in a direct sum decomposition of H_V as such a sum. Then $Z(H_V) = \Bbbk e_V$ (since $Z(\mathrm{End}_\Bbbk(V)) = \Bbbk \mathrm{id}_V$) and so $Z(H) = \bigoplus_{V\in\mathcal{V}} \Bbbk e_V$. Moreover, for any $x \in H$, we have that

$$\mathrm{Tr}(x\,\mathrm{id}_H) = \sum_{V\in\mathcal{V}} \mathrm{Tr}(x\,\mathrm{id}_{H_V}) = \sum_{V\in\mathcal{V}} \dim_\Bbbk(V)\,\mathrm{Tr}(x\,\mathrm{id}_V). \tag{4.37}$$

The map $x \in H \mapsto \mathrm{Tr}((x\,\mathrm{id}_H)\circ S^2) \in \Bbbk$ is a right integral for H^* (by [Rad94b, Proposition 2(b)] applied to H^{op}). Therefore, since $S^2 = \mathrm{id}_H$ and by the uniqueness of integrals, there exists $k \in \Bbbk$ such that $\mathrm{Tr}(x\,\mathrm{id}_H) = k\,\lambda(x)$ for all $x \in H$. Evaluating with $x = 1$, we get that $k = \dim_\Bbbk(H)/\lambda(1)$. Hence, by (4.37),

$$k\,\lambda(x) = \sum_{V\in\mathcal{V}} \dim_\Bbbk(V)\,\mathrm{Tr}(x\,\mathrm{id}_V) \tag{4.38}$$

for all $x \in H$.

Let $V \in \mathcal{V}$. By (4.21) and since $S^2 = \mathrm{id}_H$, the special grouplike element G of H is central and so $G^{-1}\mathrm{id}_V$ is H-linear. Therefore, since V is scalar and G is invertible, there exists a (unique) $\gamma_V \in \Bbbk^*$ such that $G^{-1}\mathrm{id}_V = \gamma_V \mathrm{id}_V$. Since H and H^* are semisimple and so unimodular, their special grouplike elements are trivial. Then $G^2 = 1$ by (4.22) and so $\gamma_V^2 = 1$. Hence, for all $x \in H$,

$$\begin{aligned}\dim_q(V)\,\mathrm{Tr}(G^{-1}x\,\mathrm{id}_V) &= \mathrm{Tr}(G^{-1}\mathrm{id}_V)\,\mathrm{Tr}(xG^{-1}\mathrm{id}_V)\\ &= \gamma_V^2\,\mathrm{Tr}(\mathrm{id}_V)\,\mathrm{Tr}(x\,\mathrm{id}_V) = \dim_\Bbbk(V)\,\mathrm{Tr}(x\,\mathrm{id}_V).\end{aligned} \tag{4.39}$$

For any $x \in H$, we have that:

$$\begin{aligned}\lambda(\dim_q(V)\,z_V x) &= \dim_q(V)\,\mathrm{Tr}(G^{-1}x\,\mathrm{id}_V) \quad \text{by (4.36)}\\ &= \dim_\Bbbk(V)\,\mathrm{Tr}(x\,\mathrm{id}_V) \quad \text{by (4.39)}\\ &= \textstyle\sum_{W\in\mathcal{V}} \dim_\Bbbk(W)\,\mathrm{Tr}(xe_V\,\mathrm{id}_W) \quad \text{since } e_V\,\mathrm{id}_W = \delta_{V,W}\,\mathrm{id}_V\\ &= \lambda(k\,e_V x) \quad \text{by (4.38)},\end{aligned}$$

and so $\dim_q(V)\,z_V = k\,e_V$ (since H^* is a free right H-module with basis λ).

Finally, since $k = \dim_\Bbbk(H)/\lambda(1) \neq 0$ (because the characteristic of $\Bbbk$ is 0) and $\dim_q(V) \neq 0$ (because rep_H is semisimple, see Section 3.1), we get that $Z(H) = \bigoplus_{V\in\mathcal{V}} \Bbbk e_V = \bigoplus_{V\in\mathcal{V}} \Bbbk z_V$ and $\sum_{V\in\mathcal{V}} \dim_q(V)\,z_V = k\sum_{V\in\mathcal{V}} e_V = k\,1 \in \Bbbk^* 1$. □

4.8. **HKR-type invariants.** Let H be a finite-dimensional ribbon Hopf algebra. We use notations of Section 4.6.

By Theorem 2.5 and Proposition 2.3, for any $z \in \mathcal{I}(H)^{\mathrm{norm}}$,

$$\tau_{(H,z)}(M) = \tau_{\mathrm{rep}_H}(M;\phi_z) \in \Bbbk \tag{4.40}$$

is an invariant of 3-manifolds. Note that the choice of the normalisation in the definition of $\tau_{\mathrm{rep}_H}(M;\phi_z)$ (see Proposition 2.3) implies that $\tau_{(H,z)}(M)$ does not depend on the choice of the non-zero right integral λ for H^* used to define $\tau_{\mathrm{rep}_H}(M;\phi_z)$.

By Remarks 2.6, for any $z \in \mathcal{I}(H)^{\mathrm{norm}}$, we have that

$$\tau_{(H,z)}(S^3) = 1, \qquad \tau_{(H,z)}(M\#M') = \tau_{(H,z)}(M)\,\tau_{(H,z)}(M'),$$

for all 3-manifolds M, M'. Moreover, if $z \in \mathcal{I}(H)^{\mathrm{norm}}$, $n \in N(H)$, and $k \in \Bbbk^*$, then $kz + n \in \mathcal{I}(H)^{\mathrm{norm}}$ and, for all 3-manifold M,

$$\tau_{(H,kz+n)}(M) = \tau_{(H,z)}(M). \tag{4.41}$$

Definition 4.15. An invariant of closed 3-manifolds I with values in $\Bbbk$ is said to be of *HKR-type* if there exist a finite-dimensional ribbon Hopf algebra H (over $\Bbbk$) and an element $z \in \mathcal{I}(H)^{\mathrm{norm}}$ such that $I(M) = \tau_{(H,z)}(M)$ for all 3-manifolds M.

In Proposition 4.18, we show that the Reshetikhin-Turaev invariants defined with premodular Hopf algebras are of HKR-type.

Let us show that any HKR-type invariant can be computed by using the Kauffman-Radford algorithm (which is given in [KR95] for the case H unimodular and for $z = 1$). Fix a finite-dimensional ribbon Hopf algebra H, a non-zero right integral λ for H^*, and an element $z \in \mathcal{I}(H)^{\mathrm{norm}}$. Let M be a 3-manifold and $L = L_1 \cup \cdots \cup L_n$ be a framed link in S^3 such that $M \simeq M_L$. Let us recall the Kauffman-Radford algorithm[1]:

(A). Consider a diagram D of L (with blackboard framing). Each crossing of D is decorated with the R-matrix $R = \sum_i a_i \otimes b_i$ as in Figure 11. The diagram obtained after this step is called the *flat diagram of* D. Note that the flat diagram of D is composed by n closed plane curves, each of them arising from a component of L.



FIGURE 11.

(B). On each component of the flat diagram of D, the algebraic decoration is concentrated in an arbitrary point (other than extrema and crossings) according to the rules of Figure 12, where $a, b \in H$.



FIGURE 12.

In that way we get an element $\sum_k v_1^k \otimes \cdots \otimes v_n^k \in H^{\otimes n}$, where v_i^k corresponds to the component of the flat diagram of D arising from L_i, see Figure 13.



FIGURE 13.

---

[1] This corresponds to [KR95] applied with the opposite ribbon Hopf algebra H^{op} to H.

For $1 \leq i \leq n$, let d_i be the Whitney degree of the flat diagram of L_i obtained by traversing it upwards from the point where the algebraic decoration have been concentrated. The Whitney degree is the total turn of the tangent vector to the curve when one traverses it in the given direction, see Figure 14.



FIGURE 14.

Proposition 4.16.

$$\tau_{(H,z)}(M) = \lambda(z\theta)^{b_-(L)-n_L}\, \lambda(z\theta^{-1})^{-b_-(L)} \sum_k \lambda(zG^{d_1+1}v_1^k) \cdots \lambda(zG^{d_n+1}v_n^k).$$

Proof. Choose an orientation for L. Let T be a n-special tangle such that L is isotopic $T \circ (\cup_- \otimes \cdots \otimes \cup_-)$, where the ith cup (with clockwise orientation) corresponds to the component L_i. Let D_T be a diagram of T. Note that $D = D_T \circ (\cup \otimes \cdots \otimes \cup)$ is a diagram of L. Apply steps (A) and (B) to D_T, see Figure 15. Note that, in this case, $d_i = -1$.



FIGURE 15.

From the definition of the monoidal structure, duality, braiding and twist of rep_H (see Section 4.4), it is not difficult to verify that, for any finite-dimensional left H-modules $M_1, \ldots, M_n$,

$$T_{(M_1,\ldots,M_n)} = \sum_k \mathrm{ev}_{M_1}(\mathrm{id}_{M_1^*} \otimes v_k^1\, \mathrm{id}_{M_1}) \otimes \cdots \otimes \mathrm{ev}_{M_n}(\mathrm{id}_{M_n^*} \otimes v_k^n\, \mathrm{id}_{M_n}).$$

Then, by Lemma 4.3,

$$\begin{aligned}\tau_{\mathrm{rep}_H}(L;\phi_z) = \phi_T \circ \phi_z^{\otimes n} &= \sum_k \mathrm{ev}_H(\phi_z(1_\Bbbk) \otimes v_k^1) \otimes \cdots \otimes \mathrm{ev}_H(\phi_z(1_\Bbbk) \otimes v_k^n)\\ &= \sum_k \lambda(zv_1^k) \cdots \lambda(zv_n^k) = \sum_k \lambda(zG^{d_1+1}v_1^k) \cdots \lambda(zG^{d_n+1}v_n^k).\end{aligned}$$

Hence we get the results since $\Theta_\pm \phi_z = \lambda(z\theta^{\pm 1})$. □

Corrolary 4.17. *Suppose that H is unimodular and $\lambda(\theta) \neq 0 \neq \lambda(\theta^{-1})$. Then $1 \in \mathcal{I}(H)^{\mathrm{norm}}$ and $\tau_{(H,1)}(M)$ is the Hennings-Kauffman-Radford invariant of 3-manifolds defined with the opposite ribbon Hopf algebra H^{op} to H.*

Proof. This is an immediate consequence of Corollary 4.9, Proposition 4.16, and the definition of the Hennings-Kauffman-Radford invariant given, e.g., in [KR95]. □

4.9. **Reshetikhin-Turaev from premodular Hopf algebras.** Let $(H,\mathcal{V})$ be a finite-dimensional *premodular* Hopf algebra. This means (see [Tur94]) that H is a finite-dimensional ribbon Hopf algebra and $\mathcal{V}$ is a finite set of finite-dimensional pairwise non-isomorphic left H-modules such that:

- each $V \in \mathcal{V}$ is non-negligible scalar;
- the trivial left H-module $\Bbbk$ belongs to $\mathcal{V}$;
- for any $V \in \mathcal{V}$, there exists $W \in \mathcal{V}$ such that $V^* \simeq W$;
- for any $V, W \in \mathcal{V}$, $V \otimes W$ splits as a (finite) direct sum of certain modules of $\mathcal{V}$ (possibly with multiplicities) and a negligible H-module.

By *negligible* H-module, we mean a finite-dimensional left H-module N such that $\mathrm{tr}_q(f)=0$ for any $f \in \mathrm{End}_{\mathrm{rep}_H}(N)$ or, equivalently, such that $\dim_q(N)=0$.

Consider the semisimplification rep_H^s of rep_H (see Section 3.4) and let $\pi : \mathrm{rep}_H \to \mathrm{rep}_H^s$ be the ribbon functor associated to this semisimplification.

Let $\mathcal{B}_\mathcal{V}$ be the full subcategory of rep_H^s whose objects are finite directs sums of objects of $\pi(\mathcal{V}) = \{\pi(V) \,|\, V \in \mathcal{V}\}$. By the definition of a premodular Hopf algebra, we have that $\mathcal{B}_\mathcal{V}$ is a full ribbon subcategory of rep_H^s. Note that $\mathcal{B}_\mathcal{V}$ is a finitely semisimple ribbon $\Bbbk$-category whose simples objects are scalar and has $\pi(\mathcal{V})$ as a (finite) set of representatives of isomorphism classes of simple objects. Recall that the Reshetikhin-Turaev invariant $\mathrm{RT}_{\mathcal{B}_\mathcal{V}}(M)$ of 3-manifolds is well-defined when $\Delta_\pm^{\mathcal{B}_\mathcal{V}} \neq 0$ (see Section 3.3).

Let λ be a non-zero right integral for H^*. For any $V \in \mathcal{V}$, as in (4.36), we let $z_V \in H$ such that $\lambda(z_V x) = \mathrm{Tr}(G^{-1} x \,\mathrm{id}_V)$ for all $x \in H$. Set

$$z_\mathcal{V} = \sum_{V \in \mathcal{V}} \dim_q(V)\, z_V, \tag{4.42}$$

where $\dim_q(V) = \mathrm{Tr}(G^{-1}\mathrm{id}_V)$ is the quantum dimension of V. By Corollary 4.10(b), we have that $z_\mathcal{V} \in \mathcal{I}(H)$.

Proposition 4.18. *If $\Delta_\pm^{\mathcal{B}_\mathcal{V}} \neq 0$, then $z_\mathcal{V} \in \mathcal{I}(H)^{\mathrm{norm}}$ and $\tau_{(H,z_\mathcal{V})}(M) = \mathrm{RT}_{\mathcal{B}_\mathcal{V}}(M)$ for all 3-manifold M.*

Note that Proposition 4.18 says that the Reshetikhin-Turaev invariant defined from a premodular Hopf algebra is of HKR-type.

Proof. Let (A,i) be the coend of the functor (1.16) of rep_H (as in Section 4.5), let (B,j) be the coend of the functor (1.16) of $\mathcal{B}_\mathcal{V}$ (as in Section 3.2). Set $\alpha_{\mathcal{B}_\mathcal{V}} = \sum_{V \in \mathcal{V}} \dim_q(V) j_{\pi(V)} \widetilde{\mathrm{coev}}_{\pi(V)}$. Suppose that $\Delta_\pm^{\mathcal{B}_\mathcal{V}} \neq 0$. By Corollary 3.7, we have that $\alpha_{\mathcal{B}_\mathcal{V}} \in \mathcal{I}(\mathcal{B}_\mathcal{V})^{\mathrm{norm}}$ and $\mathrm{RT}_{\mathcal{B}_\mathcal{V}}(M) = \tau_{\mathcal{B}_\mathcal{V}}(M;\alpha_{\mathcal{B}_\mathcal{V}})$ for all 3-manifold M. Set $\varphi_{\mathcal{B}_\mathcal{V}} : B \to \pi(A)$ as in (3.10). As in the proof of Corollary 4.10, we have that $\pi(\phi_{z_V}) = \varphi_{\mathcal{B}_\mathcal{V}} j_{\pi(V)} \widetilde{\mathrm{coev}}_{\pi(V)}$. Then $\pi(\phi_{z_\mathcal{V}}) = \varphi_{\mathcal{B}_\mathcal{V}} \alpha_{\mathcal{B}_\mathcal{V}}$. Since $\alpha_{\mathcal{B}_\mathcal{V}} \in \mathcal{I}(\mathcal{B}_\mathcal{V})^{\mathrm{norm}}$ and $\pi(\phi_{z_\mathcal{V}}) = \varphi \alpha_{\mathcal{B}_\mathcal{V}}$, Corollary 3.9(b) gives that $\phi_{z_\mathcal{V}} \in \mathcal{I}(\mathrm{rep}_H)^{\mathrm{norm}}$ and $\tau_{\mathcal{B}_\mathcal{V}}(M;\alpha_{\mathcal{B}_\mathcal{V}}) = \tau_{\mathrm{rep}_H}(M;\phi_{z_\mathcal{V}})$ for all 3-manifold M. Hence $z_\mathcal{V} \in \mathcal{I}(H)^{\mathrm{norm}}$ and $\tau_{(H,z_\mathcal{V})}(M) = \mathrm{RT}_{\mathcal{B}_\mathcal{V}}(M)$ for all 3-manifold M. □

Note that if H is semisimple finite-dimensional ribbon Hopf algebra, $\Bbbk$ is a splitting field for H, and $\mathcal{V}$ is a set of representatives of isomorphism classes of simple left H-modules, then $(H,\mathcal{V})$ is a premodular Hopf algebra and $\mathcal{B}_\mathcal{V} = \mathrm{rep}_H$.

Corrolary 4.19. *Let H be a finite-dimensional semisimple ribbon Hopf algebra. Suppose that the base field $\Bbbk$ is of characteristic 0 and is a splitting field for H. Then the Hennings-Kauffman-Radford invariant of 3-manifolds computed with H^{op} and* *the Reshetikhin-Turaev invariant of 3-manifolds computed with* rep_H *are simultaneously well-defined (that is,* $\Delta_{\pm}^{\operatorname{rep}_H} \neq 0$ *if and only if* $1 \in \mathcal{I}(H)^{\mathrm{norm}}$*). Moreover, if they are, then they coincide, that is,* $\tau_{(H,1)}(M) = \mathrm{RT}_{\operatorname{rep}_H}(M)$ *for all 3-manifold* M*.*

Remark 4.20. Conclusions of Corollary 4.19 may be no more true when H is not semisimple (see Remark 4.12). Moreover, in the modular case (in the sense of Remark 3.5), Corollary 4.19 was first shown in [Ker97].

Proof. By Proposition 4.18, the Reshetikhin-Turaev invariant of 3-manifolds computed from rep_H is well-defined if $z_{\mathcal{V}} = \sum_{V\in\mathcal{V}} \dim_q(V)\, z_V \in \mathcal{I}(H)^{\mathrm{norm}}$ and is equal to $\tau_{(H,z_{\mathcal{V}})}$. By Corollary 4.17, the Hennings-Kauffman-Radford invariant of 3-manifolds computed with H^{op} is well-defined if $1 \in \mathcal{I}(H)^{\mathrm{norm}}$ and is equal to $\tau_{(H,1)}$. Now, by Proposition 4.14, $\sum_{V\in\mathcal{V}} \dim_q(V)\, z_V = k\,1$ for some $k \in \Bbbk^*$. We conclude by using (4.41). □

5. A NON-UNIMODULAR EXAMPLE

Let us examine the case of family of non-unimodular ribbon Hopf algebras, defined by Radford [Rad94a], which includes Sweedler's Hopf algebra.

Let n be an odd positive integer and $\Bbbk$ be a field whose characteristic does not divide $2n$. Let H_n be the $\Bbbk$-algebra generated by a and x with the following relations

$$a^{2n} = 1, \quad x^2 = 0, \quad ax = -xa.$$

The algebra H_n is a Hopf algebra for the following structure maps:

$$\begin{aligned} \Delta(a) &= a \otimes a, & \Delta(x) &= x \otimes a^n + 1 \otimes x, \\ \varepsilon(a) &= 1, & \varepsilon(x) &= 0, \\ S(a) &= a^{-1}, & S(x) &= a^n x. \end{aligned}$$

The set $\mathcal{B} = \{a^l x^m \mid 0 \le l < 2n,\ 0 \le m \le 1\}$ is a basis for H_n. The dual basis of $\mathcal{B}$ is $\{\overline{a^k x^r} \mid 0 \le k < 2n,\ 0 \le r \le 1\}$, where $\overline{a^k x^n}(a^l x^r) = \delta_{l,k}\,\delta_{m,r}$. Set

$$\Lambda = (1 + a + a^2 + \cdots + a^{2n-1})x \quad \text{and} \quad \lambda = \overline{a^n x}.$$

Then Λ is a left integral for H_n and λ is a right integral for H_n^* such that $\lambda(\Lambda) = \lambda(S(\Lambda)) = 1$. The distinguished grouplike element of H_n is $g = a^n \in G(H_n)$ and the distinguished grouplike element $\nu \in G(H_n^*) = \mathrm{Alg}(H_n, \Bbbk)$ of H_n^* is given by $\nu(a) = -1$ and $\nu(x) = 0$.

Suppose that $\Bbbk$ has a primitive $2n$-root of unity ω. Let an odd integer $1 \le s < 2n$ and $\beta \in \Bbbk$. Then

$$R_{\omega,s,\beta} = \frac{1}{2n} \sum_{0\le i,l<2n} w^{-il}\, a^i \otimes a^{sl} + \frac{\beta}{2n} \sum_{0\le i,l<2n} w^{-il}\, a^i x \otimes a^{sl+n} x$$

is a R-matrix for H_n and $h_\nu = (\mathrm{id}_{H_n} \otimes \nu)(R_{\omega,s,\beta}) = a^n$.

The set $\{e_l = \frac{1}{2n} \sum_{0\le i<2n} \omega^{-il} a^i \mid 0 \le l < 2n\}$ is a basis of the algebra $\Bbbk[a]$ and verifies:

$$\begin{aligned} &e_i e_j = \delta_{i,j}\, e_i, && 1 = e_0 + \cdots + e_{2n-1}, && a^i e_j = \omega^{ij} e_j, \\ &\varepsilon(e_l) = \delta_{l,0}, && \Delta(e_l) = \sum_{0\le i<2n} e_i \otimes e_{l-i}, && S(e_l) = e_{-l}. \end{aligned}$$

Denote by $\chi : \Bbbk^* \to \Bbbk[a]$ the algebra homomorphism defined by

$$\alpha \in \Bbbk^* \mapsto \chi(\alpha) = \sum_{0 \leq l < 2n} \alpha^{l^2}\, e_l.$$

The quasitriangular Hopf algebra $(H_n, R_{\omega,s,\beta})$ is ribbon with twist $\theta = a^n \chi(\omega^s)$. Note that $\theta^p = a^{pn}\chi(\omega^{ps})$ for any integer p. The special grouplike element of H_n is then $G = \sum_{0 \leq l < 2n} (-1)^l\, e_l = a^n$.

Let $T : H_n \to H_n$, $L(H_n)$, $N(H_n)$, $V_2(H_n)$, $\mathcal{I}(H_n)$, and $\mathcal{I}(H_n)^{\mathrm{norm}}$ be as in Section 4.6. It is not difficult to verify that

$$\begin{aligned}
&T(a^k) = (-1)^k a^{n-k} \quad \text{and} \quad T(a^k x) = a^{-k} x \quad \text{for all } 0 \leq k < 2n,\\
&L(H_n) = \Bbbk x \oplus \Bbbk a x \oplus \cdots \oplus \Bbbk a^{2n-1} x,\\
&Z(H_n) = \Bbbk 1 \oplus \Bbbk a^2 \oplus \cdots \oplus \Bbbk a^{2n-2},\\
&N(H_n) = \Bbbk x \oplus \Bbbk a^2 x \oplus \cdots \oplus \Bbbk a^{2n-2} x = Z(H_n)x,\\
&V_2(H_n) = \underset{0 \leq p,q < n}{\oplus} \Bbbk(a^{2p} \otimes a^{2q}) \oplus \underset{0 \leq k,l < 2n}{\oplus} \Bbbk(a^k x \otimes a^l x).
\end{aligned}$$

Let $z \in L(H_n)$. Since $a^{2n} = 1$, we can write $z = \sum_{k \in \mathbb{Z}/2n\mathbb{Z}} \alpha_k a^k x$ for some function $\alpha : \mathbb{Z}/2n\mathbb{Z} \to \Bbbk$. Using Theorem 4.8, we get that $z \in \mathcal{I}(H_n)$ if and only if

$$\alpha_{-k} = \alpha_k \quad \text{and} \quad \alpha_k \alpha_{k+l-n} = \alpha_k \alpha_l.$$

for all k, l odd. Let $\gamma : \alpha : \mathbb{Z}/n\mathbb{Z} \to \Bbbk$ defined by $\gamma_k = \alpha_{2k+n}$. Since n is odd, we have that $z - \sum_{k \in \mathbb{Z}/n\mathbb{Z}} \gamma_k a^{2k+n} x \in N(H_n)$. Then $z \in \mathcal{I}(H_n)$ if and only if

$$\gamma_{-k} = \gamma_k \quad \text{and} \quad \gamma_k \gamma_{k+l} = \gamma_k \gamma_l. \tag{5.1}$$

for all k, l.

Suppose that $z \in \mathcal{I}(H_n)$ and $z \neq 0$. Using (5.1), we get that $\gamma_k \gamma_0 = \gamma_k \gamma_{-k} = \gamma_k^2$ for all k. In particular, $\gamma_0 \neq 0$ and $\gamma_k = \gamma_0$ whenever $\gamma_k \neq 0$. Set

$$d = \min\{1 \leq k \leq n \,|\, \gamma_k \neq 0\} \geq 1.$$

Note that $\gamma_k = 0$ for all $1 \leq k < d$ and, by (5.1), that $\gamma_{k+d} = \gamma_k$ for all k. The integer d divides n. Indeed, let r such that $rd \leq n < rd + d$. Then $0 \leq n - rd \leq n$ and $\gamma_{n-rd} = \gamma_n = \gamma_0 \neq 0$. Therefore, by definition of d, we get that $n - rd = 0$ and so $d|n$. Hence $z = \gamma_0 z_d + w$, where $w \in N(H_n)$ and

$$z_d = \sum_{k=0}^{\frac{n}{d}-1} a^{2dk+n} x.$$

Conversely, it is not difficult to verify that $z_d \in \mathcal{I}(H_n)$ whenever d divides n. Hence

$$\mathcal{I}(H_n) = \bigcup_{d|n} \{\alpha z_d + w \,|\, \alpha \in \Bbbk \text{ and } w \in N(H_n)\}.$$

Let d be an positive integer dividing n. For any $\alpha \in \Bbbk$ and $w \in N(H_n)$, we have that

$$\lambda((\alpha z_d + w)\theta^{\pm 1}) = \frac{\alpha}{2d} \sum_{k=0}^{2d-1} \left(\omega^{\frac{n}{d}}\right)^{\pm s \frac{n}{d} k^2 + nk}.$$

The sum of the right-hand sum of this equality is a Gauss sum which is non-zero if and only if the enhancement $k \in \mathbb{Z}/2d\mathbb{Z} \mapsto \psi(k) = \pm s\frac{n}{d}k^2 + nk \in \mathbb{Z}/2d\mathbb{Z}$ is tame, that is, $\psi(x) = 0$ for any $x \in \mathbb{Z}/2d\mathbb{Z}$ such that $\psi(x+y) = \psi(x) + \psi(y)$ for all

$y \in \mathbb{Z}/2d\mathbb{Z}$, see [Tay]. Since n and s are odd, it is not difficult to verify that ψ is tame. Therefore $\lambda((\alpha z_d + w)\theta^{\pm 1}) \neq 0$ if and only if $\alpha \neq 0$. Hence

$$\mathcal{I}(H_n)^{\mathrm{norm}} = \bigcup_{d|n} \{\alpha z_d + w \mid \alpha \in \Bbbk^* \text{ and } w \in N(H_n)\}.$$

In conclusion, the ribbon Hopf algebra H_n leads to $D(n)$ HKR-type invariants of 3-manifolds, where $D(n)$ denotes the number of positive divisors of n, which are $\tau_{(H_n, z_d)}$ where $1 \leq d \leq n$ and $d|n$.

Note that H_n is not unimodular (and so neither semisimple) since $\nu \neq \varepsilon$. Therefore $1 \notin \mathcal{I}(H_n)$ (by Corollary 4.9), that is, the Hennings-Kauffman-Radford invariant is not defined for H_n. Moreover, the categorical Hopf algebra $A = H_n^*$ of rep_{H_n} does not possess any non-zero two-sided integral (since H_n is not unimodular), and so the Lyubashenko invariant of 3-manifolds is not defined for rep_{H_n}.

Appendix A. Traces on ribbon Hopf algebras

Recall that a *trace* on a Hopf algebra H is a form $t \in H^*$ such that $t(xy) = t(yx)$ and $t(S(x)) = t(x)$ for all $x, y \in H$.

Let H be a finite-dimensional ribbon Hopf algebra. Let $\lambda \in H^*$ be a non-zero right integral for H^*, $\nu \in G(H^*) = \mathrm{Alg}(H, \Bbbk)$ be the distinguished grouplike element of H^* and G be the special grouplike element of H. Recall that $\cdot$ denotes the right action of H on H^* defined in (4.1), that $\leftharpoonup$ denotes the right H^*-action on H defined in (4.2), that $L(H)$ denotes the $\Bbbk$-subspace of H constituted by the elements $z \in H$ verifying $(x \leftharpoonup \nu)z = zx$ for all $x \in H$, and that T denotes the $\Bbbk$-endomorphism of H defined by $z \mapsto T(z) = (S(z) \leftharpoonup \nu)h_\nu$, where $h_\nu = (\mathrm{id}_H \otimes \nu)(R) \in G(H)$.

In the next proposition, we give an algebraic description of the space of traces on H.

Proposition A.1. *The space $\{z \in L(H) \mid T(z) = z\}$ is $\Bbbk$-isomorphic to the space of traces on H via the map $z \mapsto \lambda \cdot (zG)$.*

If H is unimodular, then $L(H) = Z(H)$ and $T = S$, and so we recover the parameterization of traces on H given in [Rad94b, Hen96].

Proof. Let $z \in L(H)$ such that $T(z) = z$. Set $t = \lambda \cdot (zG) \in H^*$. By using (4.26) and (4.21), we have that for any $x, y \in H$,

$$t(xy) = \lambda(zGxy) = \lambda(zS^2(y)Gx) = \lambda(zGyx) = t(yx).$$

Moreover, for any $x \in H$,

$$\begin{aligned}
t(S(x)) &= \lambda(zGS(x)) = \lambda(G^2 h_\nu x G^{-1} S^{-1}(z)) \quad \text{by (4.23)} \\
&= \lambda((S(z) \leftharpoonup \nu) G^2 h_\nu x G^{-1}) \quad \text{by (4.3)} \\
&= \lambda(z h_\nu^{-1} G^2 h_\nu x G^{-1}) \quad \text{since } T(z) = z \\
&= \lambda(z S^2(G^{-1}) h_\nu^{-1} G^2 h_\nu x) \quad \text{by (4.26)} \\
&= \lambda(z G^{-1} h_\nu^{-1} G^2 h_\nu x) \\
&= \lambda(z G S^{-4}(h_\nu^{-1}) h_\nu x) \quad \text{by (4.21)} \\
&= \lambda(z G h_\nu^{-1} h_\nu x) = \lambda(zGx) = t(x).
\end{aligned}$$

Conversely, let $t \in H^*$ be a trace on H. Since $(H^*, \cdot)$ is a free right H-module of rank 1 with basis λ and G is invertible, there exists a (unique) $z \in H$ such that $t = \lambda \cdot (zG)$. Let $x \in H$. For all $a \in H$,

$$\begin{aligned}
\lambda(zxGy) &= \lambda(zGS^{-2}(x)y) \quad \text{by (4.21)} \\
&= t(S^{-2}(x)y) = t(yS^{-2}(x)) \\
&= \lambda(zGyS^{-2}(x)) \\
&= \lambda((x \leftharpoonup \nu)zGy) \quad \text{by (4.3).}
\end{aligned}$$

Therefore, since $(H^*, \cdot)$ is a free right H-module of rank 1 with basis λ, we have that $(x \leftharpoonup \nu)zG = zxG$ and so $(x \leftharpoonup \nu)z = zx$. Hence $z \in L(H)$. Moreover, for all $x \in H$,

$$\begin{aligned}
\lambda(zGx) &= t(x) = t(S(x)) = \lambda(zGS(x)) \\
&= \lambda(G^2 h_\nu x G^{-1} S^{-1}(z)) \quad \text{by (4.23)} \\
&= \lambda(G^2 h_\nu x S^{-1}(zG)) \\
&= \lambda(G^2 h_\nu x S^{-1}((G \leftharpoonup \nu)z)) \quad \text{since } z \in L(H) \\
&= \lambda((S((G \leftharpoonup \nu)z) \leftharpoonup \nu)G^2 h_\nu x) \quad \text{by (4.3).}
\end{aligned}$$

Therefore, since $(H^*, \cdot)$ is a free right H-module of rank 1 with basis λ,

$$\begin{aligned}
zG &= (S((G \leftharpoonup \nu)z) \leftharpoonup \nu)G^2 h_\nu \\
&= (S(z) \leftharpoonup \nu)(S(G \leftharpoonup \nu) \leftharpoonup \nu)GS^2(h_\nu)G \quad \text{by (4.21)} \\
&= (S(z) \leftharpoonup \nu)\nu(G)\nu(G^{-1})G^{-1}Gh_\nu G \\
&= (S(z) \leftharpoonup \nu)h_\nu G = T(z)G
\end{aligned}$$

and so $T(z) = z$. □

References

[Abe80] E. Abe, *Hopf algebras*, Cambridge University Press, Cambridge, 1980.

[Ben98] D. J. Benson, *Representations and cohomology I*, second ed., Cambridge Studies in Advanced Mathematics, vol. 30, Cambridge University Press, Cambridge, 1998.

[BKLT00] Y. Bespalov, T. Kerler, V. Lyubashenko, and V. Turaev, *Integrals for braided hopf algebras*, J. Pure Appl. Algebra **148** (2000), no. 2, 113–164.

[Bru00a] A. Bruguières, *Catégories prémodulaires, modularisations et invariants des variétés de dimension 3*, Math. Annalen **316** (2000), no. 2, 215–236.

[Bru00b] ———, *Tresses et structure entière sur la catégorie des représentations de* SL_N *quantique*, Comm. Algebra **28** (2000), no. 4, 1989–2028.

[CR62] C. Curtis and I. Reiner, *Representation theory of finite groups and associative algebras*, Interscience, New York, 1962.

[Dri90] V. G. Drinfeld, *On almost cocommutative Hopf algebras*, Leningrad Math. J. **1** (1990), no. 2, 321–342.

[Hen96] M. A. Hennings, *Invariants of links and 3-manifolds obtained from Hopf algebras*, J. of London Math. Soc. **54** (1996), no. 3, 594–624.

[Ker97] T. Kerler, *Genealogy of non-perturbative quantum-invariants of* 3*-manifolds: the surgical family*, Geometry and physics (Aarhus, 1995), Dekker, New York, 1997, pp. 503–547.

[Kir78] R. Kirby, *A calculus of framed links in* S^3, Invent. Math. **45** (1978), no. 3, 35–56.

[KR95] L. H. Kauffman and D. E. Radford, *Invariants of 3-manifolds derived from finite dimensional Hopf algebras*, J. of Knot Theory and its Ramifications **4** (1995), no. 1, 131–162.

[Lan98] S. Mac Lane, *Categories for the working mathematician*, second ed., Springer-Verlag, New York, 1998.

[Lic97] W. B. R. Lickorish, *An introduction to knot theory*, Springer-Verlag, New York, 1997.

[LR87] R. G. Larson and D. E. Radford, *Semisimple cosemisimple Hopf algebras*, Amer. J. Math. **109** (1987), no. 1, 187–195.

[LR88] ______, *Finite dimensional cosemisimple Hopf algebras in characteristic 0 are semisimple*, Journal of Algebra **117** (1988), no. 2, 267–289.

[Lyu95a] V. Lyubashenko, *Invariants of* 3*-manifolds and projective representations of mapping class groups via quantum groups at roots of unity*, Comm. Math. Phys. **172** (1995), no. 3, 467–516.

[Lyu95b] ______, *Tangles and Hopf algebras in braided categories*, J. Pure Appl. Algebra **98** (1995), no. 3, 245–278.

[Maj93] S. Majid, *Braided groups*, J. Pure Appl. Algebra **86** (1993), no. 2, 187–221.

[Rad90] D. E. Radford, *The group of automorphisms of a semisimple Hopf algebra over a field of characteristic* 0 *is finite*, Amer. J. Math. **112** (1990), no. 2, 331–357.

[Rad92] ______, *On the antipode of a quasitriangular Hopf algebra*, Journal of Algebra **151** (1992), no. 1, 1–11.

[Rad94a] ______, *On Kauffman's knot invariants arising from finite-dimensional Hopf algebras*, Advances in Hopf algebras (Chicago, IL, 1992), Lecture Notes in Pure and Appl. Math., vol. 158, Dekker, New York, 1994, pp. 205–266.

[Rad94b] ______, *The trace function and Hopf algebras*, Journal of Algebra **163** (1994), no. 3, 583–622.

[RT91] N. Yu Reshetikhin and V. G. Turaev, *Invariants of* 3*-manifolds via link polynomials and quantum groups*, Invent. Math. **103** (1991), no. 3, 547–597.

[Tay] L. Taylor, *Gauss Sums in Algebra and Topology*, `http://www.nd.edu/~taylor`, prepuplication.

[Tur94] V. G. Turaev, *Quantum invariants of knots and* 3*-manifolds*, Walter de Gruyter & Co., Berlin, 1994.

Département des Sciences Mathématiques, Université Montpellier II, Case Courrier 051, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

E-mail address: `virelizi@math.univ-montp2.fr`